

15PH-0618

1A1

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
21 December 2000 (21.12.2000)

PCT

(10) International Publication Number  
WO 00/77204 A1

(51) International Patent Classification<sup>7</sup>: C12N 15/12,  
15/63, 5/10, C12Q 1/68, C07K 14/705, G01N 33/68,  
A01K 67/027

(21) International Application Number: PCT/US00/15723

(22) International Filing Date: 8 June 2000 (08.06.2000)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:  
09/329,515 10 June 1999 (10.06.1999) US

(63) Related by continuation (CON) or continuation-in-part  
(CIP) to earlier application:  
US 09/329,515 (CIP)  
Filed on 10 June 1999 (10.06.1999)

(71) Applicant (for all designated States except US): UNIVER-  
SITY OF IOWA RESEARCH FOUNDATION [US/US];  
100 Oakdale Campus #214 TIC, Iowa City, IA 52242-5000  
(US).

(71) Applicant and

(72) Inventor: LORENZ, Eva [US/US]; 33 Willowbridge  
Drive, Durham, NC 27707 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): SCHWARTZ,  
David, A. [US/US]; 4810 Rapid Creek Road N.E., Iowa  
City, IA 52240 (US). SCHUTTE, Brian, C. [US/US];  
3378 Lower West Branch Road, Iowa City, IA 52245 (US).

(74) Agent: VIKSNINS, Ann, S.; Schwegman, Lundberg,  
Woessner & Kluth, P.O. Box 2938, Minneapolis, MN  
55402 (US).

(81) Designated States (national): AE, AG, AL, AM, AT, AU,  
AZ, BA, BB, BG, BR, BY, CA, CH, CN, CR, CU, CZ, DE,  
DK, DM, DZ, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU,  
ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS,  
LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO,  
NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TR,  
TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZW.

(84) Designated States (regional): ARIPO patent (GH, GM,  
KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZW), Eurasian  
patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European  
patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE,  
IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG,  
CI, CM, GA, GN, GW, ML, MR, NE, SN, TD, TG).

Published:

— With international search report.

For two-letter codes and other abbreviations, refer to the "Guid-  
ance Notes on Codes and Abbreviations" appearing at the begin-  
ning of each regular issue of the PCT Gazette.

(54) Title: VARIANT TLR4 NUCLEIC ACID AND USES THEREOF

Human (aa 290)	. . .	L	A	Y	L	D	Y	L	D	D	I	I	D	L	F	N	C	L	T	N	V	. . .	
Mouse (aa 289)	. . .	L	T	Y	T	N	D	F	S	D	D	I	V	K	-	F	H	C	L	A	N	V	. . .
Rat (aa 289)	. . .	L	T	Y	I	N	H	F	S	D	D	I	Y	N	-	L	N	C	L	A	N	I	. . .
Hamster (aa 289)	. . .	F	T	Y	A	N	E	F	S	E	D	I	T	D	-	F	D	C	L	A	N	V	. . .

(57) Abstract: The invention provides methods to identify polymorphisms at the human TLR4 locus, as well as methods to identify individuals at risk of indications that increase their morbidity and mortality.

WO 00/77204 A1

## VARIANT TLR4 NUCLEIC ACID AND USES THEREOF

### 5                    Statement of Government Rights

This invention was made at least in part with a grant from the Government of the United States of America (grants ES06537, ES07498 and ES05605 from the National Institute of Environmental Sciences, grant HL62628 from the National Heart and Lung Institute, and grant RR00059 from the  
10 General Clinical Research Centers Program). The Government may have certain rights in the invention.

### Background of the Invention

Endotoxin or lipopolysaccharide (LPS), released from the cell wall of  
15 bacteria, plays a central role in a broad spectrum of human disease. The pathogenic importance of LPS in gram-negative sepsis is well established. Intravenous LPS induces all of the clinical features of gram-negative sepsis, including fever, shock, leukopenia followed by leukocytosis, and disseminated intravascular coagulation (Favorite et al., 1942). Higher concentrations of  
20 circulating levels of endotoxin have been associated with manifestations of systemic inflammatory response syndrome (Wang et al., 1995) and the development of acute respiratory distress syndrome following sepsis (Brigham et al., 1986). Inhaled endotoxin can induce airflow obstruction in naive or previously unexposed subjects (Michel et al., 1992) and is the most important  
25 occupational exposure associated with the development Schwartz et al., 1995a), and progression (Schwartz et al., 1995b), of airway disease among exposed workers. The concentration of endotoxin in the domestic setting appears to be associated with the clinical severity of asthma (Michel et al., 1996). Moreover, recent studies have shown that endotoxin is a contaminant of particulate matter  
30 in air pollution and may play a role in the pathophysiologic consequences of air pollution (Bonner et al., 1998). Thus, endotoxin is an important cause of morbidity and mortality.

The ability of the host to respond to endotoxin may play an important role in determining the severity of the physiologic and biologic response to this  
35 frequently encountered toxin. In mice, genetic differences in susceptibility to

frequently encountered toxin. In mice, genetic differences in susceptibility to LPS have been established. LPS hyporesponsiveness arose spontaneously and was first identified in the C3H/HeJ strain. This strain had an LD<sub>50</sub> for LPS at least 20 times that observed in A/HeJ mice (Sultz et al., 1968). In addition to  
5 C3H/HeJ, two other mouse strains, C57BL/10ScCR (Coutinho et al., 1978) and its progenitor strain, C57BL/10ScN, (Vogel et al., 1979), are hyporesponsive to LPS.

Moreover, several reports suggest that humans may also respond differently to LPS. A patient with recurrent bacterial infections has been  
10 reported to be refractory to the *in vivo* and *in vitro* effects of LPS (Kuhns et al., 1997). Following challenge with intravenous LPS, this patient had no systemic increase in IL-6 or G-CSF and had a minimal rise in the concentration of leukocytes, TNF- $\alpha$ , and IL-8. Inter-individual differences have also been reported in the release and synthesis of cytokines by monocytes stimulated with  
15 LPS *in vitro* (Santamaria et al., 1989).

LPS is thought to cause much of its morbidity and mortality by activating kinases (DeFranco et al., 1998) that control the function of transcription factors (nuclear factor- $\kappa$ B and AP-1) and ultimately lead to production of pro-inflammatory cytokines and co-stimulatory molecules (Wright, 1999). Several  
20 lines of evidence suggest that the toll receptor (TLR) family, and specifically TLR4 and TLR2 regulate the interaction between LPS and intracellular kinases and may serve as a proximal target to interrupt LPS signaling (Wright, 1998; Medzhitov et al., 1997). Both TLR4 and TLR2 activate signaling through NF- $\kappa$ B and AP-1 in transfected human cell lines (Medzhitov et al., 1997; Yang et al.,  
25 1998), and TLR4 mediates LPS induced signal transduction (Chow et al., 1999). CD14, a glycosylphosphatidyl inositol-linked receptor that binds LPS (Poltorak et al., 1998a) enhances LPS induced TLR2 (Yang et al., 1998) and TLR4 (Chow et al., 1998) signaling, suggesting that the toll receptors interact with CD14 to initiate the cellular response to LPS. Studies in mice indicate that 1) the TLR4  
30 gene maps to the critical region in LPS hyporesponsive mice (Poltorak et al., 1998), 2) mutations in the TLR4 gene (Poltorak et al., 1998; Qureshi et al., 1999) are found in mouse strains (C3H/HeJ and C57BL10/ScCr) that are defective in

their response to LPS, and 3) disruption of the TLR4 gene results in a LPS hyporesponsive phenotype (Hoshino et al., 1999).

Thus, there is need to determine whether the human TLR4 gene is polymorphic, and whether any particular polymorphism is associated with disease, e.g., LPS hyporesponsiveness.

### **Summary of the Invention**

The invention provides a method to identify a mammal, e.g., a human, at risk of, or having, an indication associated with altered innate immunity, e.g., to bacterial infection. The method comprises contacting an amount of DNA obtained from a human physiological sample with an amount of at least one TLR4-specific oligonucleotide under conditions effective to amplify the DNA so as to yield amplified DNA. Then it is determined whether the amplified DNA comprises a nucleotide substitution, e.g., one that results in an amino acid substitution, i.e., the TLR4 DNA of the human encodes a variant TLR4. Thus, the invention is useful to detect polymorphisms in the TLR4 gene.

Normal healthy, non-asthmatic subjects demonstrate a reproducible airway response to an incremental LPS inhalation challenge test, with some subjects developing airflow obstruction when challenged with low concentrations of LPS and others virtually unaffected by high concentrations of inhaled LPS. These findings suggest that the spectrum of LPS responsiveness in humans is quite variable from one individual to the next (but reproducible within an individual), and that a substantial portion of the population may be hyporesponsive to inhaled LPS. As described hereinbelow, an incremental LPS inhalation challenge test was employed to reliably phenotype individuals as either responsive (at least a 20 % decline in the forced expiratory volume in one second ( $FEV_1$ ) after inhaling up to 41.5  $\mu$ g LPS) or hyporesponsive ( $FEV_1 > 80$  % of their baseline after inhaling 41.5  $\mu$ g of LPS) to inhaled LPS. Fifty-two (63 %) of these individuals were responsive to inhaled LPS and 31 (37 %) were hyporesponsive to inhaled LPS.

These results were employed to determine the relationship between polymorphisms in the TLR4 gene and the airway response to inhaled LPS in the 83 normal healthy, non-asthmatic subjects. Using single stranded

conformational variant (SSCV) analysis and direct sequencing, a missense mutation (A896G) was identified in the fourth exon of the TLR4 gene that results in replacement of a conserved aspartic acid residue with glycine at position 299 in the extracellular domain of the TLR4 receptor. The Asp299Gly sequence variant occurred in 3 LPS responsive (5.8 %) and 7 LPS hyporesponsive (22.6 %) study subjects ( $p=0.03$ ). Among the subjects with the common TLR4 allele ( $N=73$ ), the dose-response slope (percent decline  $FEV_1$ /cumulative dose of inhaled LPS) averaged a 1.86 % decline in  $FEV_1/\mu g$  inhaled LPS (range 0.01 %-19.78 %), while the dose-response slope for the subjects with the Asp299Gly allele ( $N=10$ ) was significantly less ( $p=0.007$ ), averaging 0.59 % decline in  $FEV_1/\mu g$  inhaled LPS (range 0.00 %-1.59 %). Thus, a sequence polymorphism in the TLR4 gene, i.e., a missense mutation (Asp299Gly) in the fourth exon of the TLR4 gene, occurs in a substantial portion of the population, and is associated with an airway hyporesponsive in humans challenged with inhaled LPS. The allelic frequency of the A896G substitution was 6.6 % in the study population, 7.9 % in a normal control population from Iowa (Lidral et al., 1998), and 3.3 % in the parental chromosomes of the CEPH population (NIH-CEPH, 1992). As also described herein, the presence of a TLR4 mutation was associated with gram negative sepsis, severity of sepsis, pre-term delivery, and respiratory distress syndrome in pre-term infants.

The invention also provides an isolated and purified nucleic acid molecule comprising a nucleic acid segment, e.g., genomic DNA or cDNA, encoding TLR4, such as a variant TLR4. Also provided are primers, oligonucleotides and probes comprising the isolated nucleic acid sequences of the invention. The nucleic acid molecules of the invention may be single stranded or double stranded.

Transfection of CHO cells with either the wild-type or the mutant (Asp299Gly) allele of the TLR4 gene demonstrated that this mutation interrupts TLR4-mediated LPS signaling. Moreover, the wild-type allele of TLR4 rescues the LPS hyporesponsive phenotype in either airway epithelial cells or alveolar macrophages obtained from individuals with the TLR4 mutation. Thus, these results provide the first genetic evidence that a common mutation causes

differences in LPS responsiveness that may contribute to several disease states in humans.

Therefore, the invention further provides an expression cassette comprising a nucleic acid molecule of the invention, a host cell transformed with the expression cassette, and TLR4 polypeptides isolated therefrom. The transformed host cells, or isolated TLR4 polypeptides, may be useful in identifying agents that modulate, i.e., enhance or inhibit, TLR4 activity. For example, an expression cassette comprising a nucleic acid molecule of the invention which encodes a variant TLR4 polypeptide is introduced to murine cells, e.g., oocytes via microinjection (see Sigmund et al., 1993). The resulting pups are screened for the presence of the nucleic acid molecule. Hence, the invention also provides a transgenic mouse, the genome of the cells of which is augmented with variant human TLR4 DNA. Human TLR4 transgenic mice of the invention have altered innate immunity, e.g., they are more susceptible to gram negative sepsis than their nontransgenic counterparts.

The invention also provides a method to treat an individual at risk of, or having, an indication associated with altered innate immunity, in which an agent that alters TLR4 activity is administered to the individual.

## **Brief Description of the Figures**

Figure 1. Genomic structure for the human TLR4 gene (exons are represented by boxes and introns are represented by lines). The coding sequence is in black and the nucleotides encoding the transmembrane domain (TM) are shaded. The positions of the introns in the published TLR4 cDNA sequence (Rock et al., 1998; Genbank Accession No. U88880) are listed. The exon (caps) and intron (lower case) sequences at each of the splice junctions is shown (sequences at 5' and 3' splice junctions for intron 1, SEQ ID NO:1 and SEQ ID NO:63, respectively; for intron 2, SEQ ID NO:2 and SEQ ID NO:64, respectively; and for intron 3, SEQ ID NO:3 and SEQ ID NO:65, respectively).

Figure 2. The human TLR4 gene is alternatively spliced. Photographs are of agarose gels containing the products amplified from a multi-tissue cDNA panel from human adult (panel a; Clontech #K1420-1, Palo Alto, CA), and human fetal (panel b; Clontech #K1425-1, Palo Alto, CA) cDNA. The tissues

included brain (B), heart (H), kidney (K), liver (Li), lung (Lu), pancreas (Pa), placenta (PI), skeletal muscle (Sk), spleen (Sp), and thymus (T). The first lane of each gel included a 100 bp molecular weight standard (MW). The forward and reverse PCR primers were derived from exons 1 and 4, respectively (see panel c) and amplified three products of 453, 333, and 167 bp. The DNA sequence for these bands showed that both exons 2 and 3 were present in the 453 bp band, that exon 2 was absent in the 333 bp product, and that both exons 2 and 3 were absent in the 167 bp band. The 453 bp and 333 bp sequences are identical to previously published sequences for the human TLR cDNA (Medzhitov et al., 1997; Rock et al., 1998). The open, closed, and shaded boxes indicate the untranslated, translated, and transmembrane domain portions of the TLR4 exons, respectively. The ends of the cDNAs were arbitrarily terminated at the stop codon (STP).

Figure 3. Mutations in the human TLR4 gene. The SSCV and sequence analysis were performed blinded to the LPS response phenotype of the study subjects. The SSCV gel in panel a contains the products amplified from two samples that are homozygous for the 1060A allele (lanes 1 and 3), a heterozygous sample (lane 2) with both the 896A and 896G alleles, and a homozygous sample (lane 4) with only the 896G allele. The SSCV gel in panel b contains the products amplified from a sample that is homozygous for the T allele at position -11 and the T allele at position 315 (lane 1), and from a sample that is heterozygous with a T and a deletion at position -11 and a T and a C allele at position 315. The nucleotide numbers are based on the previously published TLR4 cDNA sequence (Rock et al., 1998).

Figure 4. The aspartic acid residue at position 299 is conserved. A portion of the predicted amino acid sequence is aligned for the TLR4 genes from human (Rock et al., 1998; SEQ ID NO:4), mouse (Poitorak et al., 1998; SEQ ID NO:5), rat (Genbank Accession No. AF057025; SEQ ID NO:6), and hamster (D. Golenbock, SEQ ID NO:7). The position of the first amino acid in each sequence is given. The aspartic acid at position 299 is indicated with an arrow. Conserved amino acids are shaded.

Figure 5. Frequency distribution histogram of the dose-response slope (% decline in FEV<sub>1</sub>/μg LPS). The % decline in FEV<sub>1</sub>/μg LPS was calculated

following administration of the cumulative LPS dose that either resulted in at least a 20 % decline in FEV<sub>1</sub> or the decline in FEV<sub>1</sub> following a cumulative inhaled dose of 41.5 µg LPS. Subjects above the x-axis (solid bars) are homozygous for the 896A allele (WT/WT); subjects below the x-axis (open bars) are either heterozygous or homozygous (\*) for the missense Asp299Gly allele. The data is replotted in the inset after the values on the x-axis were log normalized. P values are presented for the comparison of the % decline in FEV<sub>1</sub>/µg LPS between subjects with the WT/WT genotype (N=73) and those with at least one Asp299Gly allele (N=10) using absolute values (P=0.037) and log normalized values (P=0.026). Since the distribution of the dose-response slope (% decline FEV<sub>1</sub>/cumulative dose of inhaled LPS) was highly skewed, the two-sample Monte-Carlo permutation test based on 10,000 permutations was used to calculate P values (Fisher et al., 1993) the missense mutation (N=10) using absolute values (p=0.04) and the log normalized values (p=0.04) and the log normalized values (p=0.025).

Figure 6. A) CHO/CD-14 cells that express the CD-14 receptor (generous gift of Doug Golenbock) were transfected with a wild-type or mutant TLR4 allele, and then exposed to LPS. Cells were co-transfected with a luciferase reporter construct. B) IL-1α levels from airway epithelial cells of individuals which had been genotyped for TLR4. IL-1α levels before and after LPS exposure are shown. C) IL-1α levels from airway epithelial cells from a TLR4 heterozygote (WT/Asp299Gly). The cells were transduced with a recombinant adenovirus which expresses TLR4 (Rock et al., 1998), or a recombinant adenovirus which expresses green fluorescent protein (GFP), and then exposed to LPS. D) TNF-α levels from alveolar macrophage from a TLR4 homozygote (Asp299Gly/Asp299Gly). The cells were transduced with a recombinant adenovirus which expresses TLR4 (Rock et al., 1998), and then exposed to LPS.

Figure 7. Codons.

Figure 8. Exemplary amino acid substitutions.

Figure 9. A) A partial nucleotide sequence of genomic human TLR4 DNA (SEQ ID NO:62). B) The 5' UTR, exon 1 (nt 1100-1283), and a partial sequence of intron 1 (Genbank Accession No. AF172169; SEQ ID NO:70). C) Exon 2 (nt 192-311), exon 3 (nt 556-722), and intron 2 (244 bp) (Genbank



Accession No. AF172170; SEQ ID NO:71). D) Exon 4 (nt 1691-6172) and the 3' UTR (Genbank Accession No. AF172171; SEQ ID NO:72).

### **Detailed Description of the Invention**

5           As used herein, the terms "isolated and/or purified" refer to *in vitro* preparation, isolation and/or purification of a nucleic acid molecule or polypeptide, so that it is not associated with *in vivo* substances. Thus, with respect to an "isolated nucleic acid molecule", which includes a polynucleotide of genomic, cDNA, or synthetic origin or some combination thereof, the

10 "isolated nucleic acid molecule" (1) is not associated with all or a portion of a polynucleotide in which the "isolated nucleic acid molecule" is found in nature, (2) is operably linked to a polynucleotide which it is not linked to in nature, or (3) does not occur in nature as part of a larger sequence. An isolated nucleic acid molecule means a polymeric form of nucleotides of at least 10 bases in length,

15 either ribonucleotides or deoxynucleotides or a modified form of either type of nucleotide. The term includes single and double stranded forms of DNA. The term "oligonucleotide" referred to herein includes naturally occurring, and modified nucleotides linked together by naturally occurring, and non-naturally occurring oligonucleotide linkages. Oligonucleotides are a polynucleotide

20 subset with 200 bases or fewer in length. Preferably, oligonucleotides are 10 to 60 bases in length and most preferably 12, 13, 14, 15, 16, 17, 18, 19, or 20 to 40 bases in length. Oligonucleotides are usually single stranded, e.g., for primers or probes; although oligonucleotides may be double stranded, e.g., for use in the construction of a variant. Oligonucleotides of the invention can be either sense

25 or antisense oligonucleotides. The term "naturally occurring nucleotides" referred to herein includes deoxyribonucleotides and ribonucleotides. The term "modified nucleotides" referred to herein includes nucleotides with modified or substituted sugar groups and the like. The term "oligonucleotide linkages" referred to herein includes oligonucleotides linkages such as phosphorothioate,

30 phosphorodithioate, phosphoroselenoate, phosphorodiselenoate, phosphoroanilothioate, phosphoraniladate, phosphoroamidate, and the like. An oligonucleotide can include a label for detection, if desired.

The term "isolated polypeptide" means a polypeptide encoded by genomic DNA, cDNA or recombinant RNA, or is synthetic origin, or some combination thereof, which isolated polypeptide (1) is not associated with proteins found in nature, (2) is free of other proteins from the same source, e.g., free of human proteins, (3) is expressed by a cell from a different species, or (4) does not occur in nature.

The term "sequence homology" means the proportion of base matches between two nucleic acid sequences or the proportion amino acid matches between two amino acid sequences. When sequence homology is expressed as a percentage, e.g., 50 %, the percentage denotes the proportion of matches over the length of sequence from one TLR4 allele that is compared to another TLR4 allele. Gaps (in either of the two sequences) are permitted to maximize matching; gap lengths of 15 bases or less are usually used, 6 bases or less are preferred with 2 bases or less more preferred. When using oligonucleotides as probes or treatments, the sequence homology between the target nucleic acid and the oligonucleotide sequence is generally not less than 17 target base matches out of 20 possible oligonucleotide base pair matches (85 %); preferably not less than 9 matches out of 10 possible base pair matches (90 %), and more preferably not less than 19 matches out of 20 possible base pair matches (95 %).

The term "selectively hybridize" means to detectably and specifically bind. Polynucleotides, oligonucleotides and fragments of the invention selectively hybridize to nucleic acid strands under hybridization and wash conditions that minimize appreciable amounts of detectable binding to nonspecific nucleic acids. High stringency conditions can be used to achieve selective hybridization conditions as known in the art and discussed herein. Generally, the nucleic acid sequence homology between the polynucleotides, oligonucleotides, and fragments of the invention and a nucleic acid sequence of interest is at least 65 %, and more typically with preferably increasing homologies of at least about 70 %, about 90 %, about 95 %, about 98 %, and 100 %.

Two amino acid sequences are homologous if there is a partial or complete identity between their sequences. For example, 85 % homology means that 85 % of the amino acids are identical when the two sequences are aligned

for maximum matching. Gaps (in either of the two sequences being matched) are allowed in maximizing matching; gap lengths of 5 or less are preferred with 2 or less being more preferred. Alternatively and preferably, two protein sequences (or polypeptide sequences derived from them of at least 30 amino acids in length) are homologous, as this term is used herein, if they have an alignment score of at more than 5 (in standard deviation units) using the program ALIGN with the mutation data matrix and a gap penalty of 6 or greater. See Dayhoff, M. O., in Atlas of Protein Sequence and Structure, 1972, volume 5, National Biomedical Research Foundation, pp. 101-110, and Supplement 2 to this volume, pp. 1-10. The two sequences or parts thereof are more preferably homologous if their amino acids are greater than or equal to 50 % identical when optimally aligned using the ALIGN program.

The term "corresponds to" is used herein to mean that a polynucleotide sequence is homologous (i.e., is identical, not strictly evolutionarily related) to all or a portion of a reference polynucleotide sequence, or that a polypeptide sequence is identical to a reference polypeptide sequence. In contradistinction, the term "complementary to" is used herein to mean that the complementary sequence is homologous to all or a portion of a reference polynucleotide sequence. For illustration, the nucleotide sequence "TATAC" corresponds to a reference sequence "TATAC" and is complementary to a reference sequence "GTATA".

The following terms are used to describe the sequence relationships between two or more polynucleotides: "reference sequence", "comparison window", "sequence identity", "percentage of sequence identity", and "substantial identity". A "reference sequence" is a defined sequence used as a basis for a sequence comparison; a reference sequence may be a subset of a larger sequence, for example, as a segment of a full-length cDNA or gene sequence given in a sequence listing, or may comprise a complete cDNA or gene sequence. Generally, a reference sequence is at least 20 nucleotides in length, frequently at least 25 nucleotides in length, and often at least 50 nucleotides in length. Since two polynucleotides may each (1) comprise a sequence (i.e., a portion of the complete polynucleotide sequence) that is similar between the two polynucleotides, and (2) may further comprise a sequence that is divergent

between the two polynucleotides, sequence comparisons between two (or more) polynucleotides are typically performed by comparing sequences of the two polynucleotides over a "comparison window" to identify and compare local regions of sequence similarity.

5           A "comparison window", as used herein, refers to a conceptual segment of at least 20 contiguous nucleotides and wherein the portion of the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) of 20 percent or less as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the  
10 two sequences. Optimal alignment of sequences for aligning a comparison window may be conducted by the local homology algorithm of Smith and Waterman (1981) Adv. Appl. Math. 2: 482, by the homology alignment algorithm of Needleman and Wunsch (1970) J. Mol. Biol. 48: 443, by the search for similarity method of Pearson and Lipman (1988) Proc. Natl. Acad. Sci. (U.S.A.) 85: 2444, by computerized implementations of these algorithms (GAP, BESTFIT, FASTA, and TFASTA in the Wisconsin Genetics Software Package Release 7.0, Genetics Computer Group, 575 Science Dr., Madison, Wis.), or by inspection, and the best alignment (i.e., resulting in the highest percentage of  
15 homology over the comparison window) generated by the various methods is selected.  
20

          The term "sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis) over the window of comparison. The term "percentage of sequence identity" means that two polynucleotide sequences are identical (i.e., on a nucleotide-by-nucleotide basis)  
25 over the window of comparison. The term "percentage of sequence identity" is calculated by comparing two optimally aligned sequences over the window of comparison, determining the number of positions at which the identical nucleic acid base (e.g., A, T, C, G, U, or I) occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total  
30 number of positions in the window of comparison (i.e., the window size), and multiplying the result by 100 to yield the percentage of sequence identity. The terms "substantial identity" as used herein denote a characteristic of a polynucleotide sequence, wherein the polynucleotide comprises a sequence that

has at least 85 percent sequence identity, preferably at least 90 to 95 percent sequence identity, more usually at least 99 percent sequence identity as compared to a reference sequence over a comparison window of at least 20 nucleotide positions, frequently over a window of at least 20-50 nucleotides, wherein the percentage of sequence identity is calculated by comparing the reference sequence to the polynucleotide sequence which may include deletions or additions which total 20 percent or less of the reference sequence over the window of comparison.

As applied to polypeptides, the term "substantial identity" means that two peptide sequences, when optimally aligned, such as by the programs GAP or BESTFIT using default gap weights, share at least about 80 percent sequence identity, preferably at least about 90 percent sequence identity, more preferably at least about 95 percent sequence identity, and most preferably at least about 99 percent sequence identity.

As used herein, "substantially pure" means an object species is the predominant species present (i.e., on a molar basis it is more abundant than any other individual species in the composition), and preferably a substantially purified fraction is a composition wherein the object species comprises at least about 50 percent (on a molar basis) of all macromolecular species present.

Generally, a substantially pure composition will comprise more than about 80 percent of all macromolecular species present in the composition, more preferably more than about 85 %, about 90 %, about 95 %, and about 99 %. Most preferably, the object species is purified to essential homogeneity (contaminant species cannot be detected in the composition by conventional detection methods) wherein the composition consists essentially of a single macromolecular species.

An isolated "variant" TLR4 polypeptide has at least 50 %, preferably at least about 80 %, and more preferably at least about 90 %, but less than 100 %, contiguous amino acid sequence homology or identity to the amino acid sequence of a reference (wild-type) TLR4 polypeptide. Preferably, the TLR4 polypeptides of the invention are biologically active. Biologically active polypeptides include those that induce an immune response when administered to an organism, are bound by antibodies specific for TLR4, activate signaling

through NF-KB and AP-1, interact with CD14, or induce cytokine release following LPS stimulation. While it is preferred that a variant TLR4 has at least about 0.1%, preferably at least about 1%, and more preferably at least about 10%, of the activity of wild-type TLR4, the invention includes variant TLR4 polypeptides having no detectable biological activity. Likewise, a "variant" TLR4 nucleic acid molecule has at least about 80%, preferably at least about 90% and more preferably at least about 95%, but less than 100% contiguous nucleic acid sequence homology or identity to the nucleic acid sequence of a wild-type TLR4 gene.

As used herein, an "indication or condition associated with aberrant, modified or altered innate immunity" includes, but is not limited to, hyporesponsiveness to LPS, susceptibility to infection with gram-negative bacteria, susceptibility to sepsis by gram-negative bacteria, susceptibility to chronic airway disease, susceptibility to asthma, susceptibility to arthritis, susceptibility to pyelonephritis, susceptibility to gall bladder disease, susceptibility to pneumonia, susceptibility to bronchitis, susceptibility to chronic obstructive pulmonary disease, severity of cystic fibrosis, and susceptibility to local and systemic inflammatory conditions, e.g., systemic inflammatory response syndrome (SIRS), local gram negative bacterial infection, or acute respiratory distress syndrome (ARDS).

#### A. Nucleic Acid Molecules of the Invention

##### 1. Sources of the Nucleic Acid Molecules of the Invention

Sources of nucleotide sequences from which the present nucleic acid molecules encoding TLR4, a portion (fragment) thereof, a variant thereof or the nucleic acid complement thereof, include total or polyA<sup>+</sup> RNA from any mammalian, preferably human, cellular source from which cDNAs can be derived by methods known in the art. Other sources of the DNA molecules of the invention include genomic libraries derived from any mammalian cellular source. Moreover, the present DNA molecules may be prepared *in vitro*, e.g., by synthesizing an oligonucleotide of about 100, preferably about 75, more preferably about 50, and even more preferably about 40, nucleotides in length, or by subcloning a portion of a DNA segment that encodes a particular TLR4.

## 2. Isolation of a Gene Encoding TLR4

A nucleic acid molecule encoding TLR4 can be identified and isolated using standard methods, as described by Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor, NY (1989). For example, reverse-transcriptase PCR (RT-PCR) can be employed to isolate and clone TLR4 cDNAs. Oligo-dT can be employed as a primer in a reverse transcriptase reaction to prepare first-strand cDNAs from isolated RNA which contains RNA sequences of interest, e.g., total RNA isolated from human tissue. RNA can be isolated by methods known to the art, e.g., using TRIZOL™ reagent (GIBCO-BRL/Life Technologies, Gaithersburg, MD). Resultant first-strand cDNAs are then amplified in PCR reactions.

"Polymerase chain reaction" or "PCR" refers to a procedure or technique in which amounts of a preselected fragment of nucleic acid, RNA and/or DNA, are amplified as described in U.S. Patent No. 4,683,195. Generally, sequence information from the ends of the region of interest or beyond is employed to design oligonucleotide primers comprising at least 7-8 nucleotides. These primers will be identical or similar in sequence to opposite strands of the template to be amplified. PCR can be used to amplify specific RNA sequences, specific DNA sequences from total genomic DNA, and cDNA transcribed from total cellular RNA, bacteriophage or plasmid sequences, and the like. See generally Mullis et al., Cold Spring Harbor Symp. Quant. Biol., 51, 263 (1987); Erlich, ed., PCR Technology, (Stockton Press, NY, 1989). Thus, PCR-based cloning approaches rely upon conserved sequences deduced from alignments of related gene or polypeptide sequences.

Primers are made to correspond to highly conserved regions of polypeptides or nucleotide sequences which were identified and compared to generate the primers, e.g., by a sequence comparison of other mammalian TLR4. One primer is prepared which is predicted to anneal to the antisense strand, and another primer prepared which is predicted to anneal to the sense strand, of a DNA molecule which encodes TLR4.

The products of each PCR reaction are separated via an agarose gel and all consistently amplified products are gel-purified and cloned directly into a suitable vector, such as a known plasmid vector. The resultant plasmids are

subjected to restriction endonuclease and dideoxy sequencing of double-stranded plasmid DNAs.

Another approach to identify, isolate and clone cDNAs which encode TLR4 is to screen a cDNA or genomic library. Screening for DNA fragments that encode all or a portion of a DNA encoding TLR4 can be accomplished by probing the library with a probe which has sequences that are highly conserved between genes believed to be related to TLR4, e.g., the homolog of a particular TLR4 from a different species, or by screening of plaques for binding to antibodies that specifically recognize TLR4. DNA fragments that bind to a probe having sequences which are related to TLR4, or which are immunoreactive with antibodies to TLR4, can be subcloned into a suitable vector and sequenced and/or used as probes to identify other cDNAs encoding all or a portion of TLR4.

Thus, "isolated and/or purified TLR4" nucleic acid refers to *in vitro* isolation of a nucleic acid molecule from its natural cellular environment, and from association with other components of the cell, such as nucleic acid or polypeptide, so that it can be sequenced, replicated, and/or expressed. For example, "isolated TLR4 nucleic acid" is RNA or DNA containing greater than 9, preferably 36, and more preferably 45 or more, sequential nucleotide bases that encode at least a portion of TLR4, a variant thereof, RNA or DNA complementary thereto, or which hybridizes to, RNA or DNA comprising TLR4 sequences, and remains stably bound under stringent conditions, as defined by methods well known in the art, e.g., in Sambrook et al., *supra*. Thus, the RNA or DNA is "isolated" in that it is free from at least one contaminating nucleic acid with which it is normally associated in the natural source of the RNA or DNA and is preferably substantially free of any other mammalian RNA or DNA. The phrase "free from at least one contaminating source nucleic acid with which it is normally associated" includes the case where the nucleic acid is reintroduced into the source or natural cell but is in a different chromosomal location or is otherwise flanked by nucleic acid sequences not normally found in the source cell.

As used herein, the term "recombinant nucleic acid", e.g., "recombinant DNA sequence or segment", refers to a nucleic acid, e.g., to DNA, that has been



derived or isolated from any appropriate tissue source, that may be subsequently chemically altered *in vitro*, so that its sequence is not naturally occurring, or corresponds to naturally occurring sequences that are not positioned as they would be positioned in a genome which has not been transformed with  
5 exogenous DNA. An example of DNA "derived" from a source, would be a DNA sequence that is identified as a useful fragment within a given organism, and which is then chemically synthesized in essentially pure form. An example of such DNA "isolated" from a source would be a useful DNA sequence that is excised or removed from said source by chemical means, e.g., by the use of  
10 restriction endonucleases, so that it can be further manipulated, e.g., amplified, for use in the invention, by the methodology of genetic engineering.

Thus, recovery or isolation of a given fragment of DNA from a restriction digest can employ separation of the digest on polyacrylamide or agarose gel by electrophoresis, identification of the fragment of interest by comparison of its  
15 mobility versus that of marker DNA fragments of known molecular weight, removal of the gel section containing the desired fragment, and separation of the gel from DNA. See Lawn et al., Nucleic Acids Res., 9, 6103 (1981), and Goeddel et al., Nucleic Acids Res., 8, 4057 (1980). Therefore, "isolated DNA" includes completely synthetic DNA sequences, semi-synthetic DNA sequences,  
20 DNA sequences isolated from biological sources, and DNA sequences derived from RNA, as well as mixtures thereof.

As used herein, the term "derived" with respect to a RNA molecule means that the RNA molecule has complementary sequence identity to a particular DNA molecule.

### 25 3. Variants of the Nucleic Acid Molecules of the Invention

Nucleic acid molecules encoding amino acid sequence variants of TLR4 are prepared by a variety of methods known in the art. These methods include, but are not limited to, isolation from a natural source (in the case of naturally occurring amino acid sequence variants) or preparation by oligonucleotide-  
30 mediated (or site-directed) mutagenesis, PCR mutagenesis, and cassette mutagenesis of an earlier prepared variant or a non-variant version of TLR4 nucleic acid.

Oligonucleotide-mediated mutagenesis is a preferred method for preparing amino acid substitution variants of TLR4. This technique is well known in the art as described by Adelman et al., *DNA*, 2, 183 (1983). Briefly, TLR4 DNA is altered by hybridizing an oligonucleotide encoding the desired mutation to a DNA template, where the template is the single-stranded form of a plasmid or bacteriophage containing the unaltered or native DNA sequence of TLR4. After hybridization, a DNA polymerase is used to synthesize an entire second complementary strand of the template that will thus incorporate the oligonucleotide primer, and will code for the selected alteration in the TLR4 DNA.

Generally, oligonucleotides of at least 25 nucleotides in length are used. An optimal oligonucleotide will have 12 to 15 nucleotides that are completely complementary to the template on either side of the nucleotide(s) coding for the mutation. This ensures that the oligonucleotide will hybridize properly to the single-stranded DNA template molecule. The oligonucleotides are readily synthesized using techniques known in the art such as that described by Crea et al., *Proc. Natl. Acad. Sci. U.S.A.*, 75, 5765 (1978).

The DNA template can be generated by those vectors that are either derived from bacteriophage M13 vectors (the commercially available M13mp18 and M13mp19 vectors are suitable), or those vectors that contain a single-stranded phage origin of replication as described by Viera et al., *Meth. Enzymol.*, 153, 3 (1987). Thus, the DNA that is to be mutated may be inserted into one of these vectors to generate single-stranded template. Production of the single-stranded template is described in Sections 4.21-4.41 of Sambrook et al., *Molecular Cloning: A Laboratory Manual* (Cold Spring Harbor Laboratory Press, N.Y. 1989).

Alternatively, single-stranded DNA template may be generated by denaturing double-stranded plasmid (or other) DNA using standard techniques.

For alteration of the native DNA sequence (to generate amino acid sequence variants, for example), the oligonucleotide is hybridized to the single-stranded template under suitable hybridization conditions. A DNA polymerizing enzyme, usually the Klenow fragment of DNA polymerase I, is then added to synthesize the complementary strand of the template using the oligonucleotide as

a primer for synthesis. A heteroduplex molecule is thus formed such that one strand of DNA encodes the mutated form of TLR4, and the other strand (the original template) encodes the native, unaltered sequence of TLR4. This heteroduplex molecule is then transformed into a suitable host cell, usually a prokaryote such as *E. coli* JM101. After the cells are grown, they are plated onto agarose plates and screened using the oligonucleotide primer radiolabeled with 32-phosphate to identify the bacterial colonies that contain the mutated DNA. The mutated region is then removed and placed in an appropriate vector for peptide or polypeptide production, generally an expression vector of the type typically employed for transformation of an appropriate host.

The method described immediately above may be modified such that a homoduplex molecule is created wherein both strands of the plasmid contain the mutations(s). The modifications are as follows: The single-stranded oligonucleotide is annealed to the single-stranded template as described above. A mixture of three deoxyribonucleotides, deoxyriboadenosine (dATP), deoxyriboguanosine (dGTP), and deoxyribothymidine (dTTP), is combined with a modified thiodeoxyribocytosine called dCTP-( $\alpha$ S) (which can be obtained from the Amersham Corporation). This mixture is added to the template-oligonucleotide complex. Upon addition of DNA polymerase to this mixture, a strand of DNA identical to the template except for the mutated bases is generated. In addition, this new strand of DNA will contain dCTP-( $\alpha$ S) instead of dCTP, which serves to protect it from restriction endonuclease digestion.

After the template strand of the double-stranded heteroduplex is nicked with an appropriate restriction enzyme, the template strand can be digested with ExoIII nuclease or another appropriate nuclease past the region that contains the site(s) to be mutagenized. The reaction is then stopped to leave a molecule that is only partially single-stranded. A complete double-stranded DNA homoduplex is then formed using DNA polymerase in the presence of all four deoxyribonucleotide triphosphates, ATP, and DNA ligase. This homoduplex molecule can then be transformed into a suitable host cell such as *E. coli* JM101.

For example, a preferred embodiment of the invention is an isolated and purified DNA molecule comprising a human DNA segment encoding a variant TLR4 having a nucleotide substitution at position 896 (A896G) which encodes

an amino acid substitution (Asp299Gly). Other nucleotide substitutions which result in silent mutations, missense mutations, or a nonsense mutations, can be ascertained by reference to Figure 7, Figure 8 and page D1 in Appendix D in Sambrook et al., Molecular Cloning: A Laboratory Manual (1989).

## 5 B. Chimeric Expression Cassettes

To prepare expression cassettes for transformation herein, the recombinant DNA sequence or segment may be circular or linear, double-stranded or single-stranded. A DNA sequence which encodes an RNA sequence that is substantially complementary to a mRNA sequence encoding TLR4 is typically a "sense" DNA sequence cloned into a cassette in the opposite  
10 orientation (i.e., 3' to 5' rather than 5' to 3'). Generally, the DNA sequence or segment is in the form of chimeric DNA, such as plasmid DNA, that can also contain coding regions flanked by control sequences which promote the expression of the DNA present in the resultant cell line.

15 As used herein, "chimeric" means that a vector comprises DNA from at least two different species, or comprises DNA from the same species, which is linked or associated in a manner which does not occur in the "native" or wild type of the species.

Aside from DNA sequences that serve as transcription units for TLR4, or  
20 portions thereof, a portion of the recombinant DNA may be untranscribed, serving a regulatory or a structural function. For example, the DNA may itself comprise a promoter that is active in mammalian cells, or may utilize a promoter already present in the genome that is the transformation target. Such promoters include the CMV promoter, as well as the SV40 late promoter and retroviral  
25 LTRs (long terminal repeat elements), although many other promoter elements well known to the art may be employed in the practice of the invention.

Other elements functional in the host cells, such as introns, enhancers, polyadenylation sequences and the like, may also be a part of the DNA. Such elements may or may not be necessary for the function of the DNA, but may  
30 provide improved expression of the DNA by affecting transcription, stability of the mRNA, or the like. Such elements may be included in the DNA as desired to obtain the optimal performance of the transforming DNA in the cell.

"Control sequences" is defined to mean DNA sequences necessary for the expression of an operably linked coding sequence in a particular host organism. The control sequences that are suitable for prokaryotic cells, for example, include a promoter, and optionally an operator sequence, and a ribosome binding site. Eukaryotic cells are known to utilize promoters, polyadenylation signals, and enhancers.

"Operably linked" is defined to mean that the nucleic acids are placed in a functional relationship with another nucleic acid sequence. For example, DNA for a presequence or secretory leader is operably linked to DNA for a peptide or polypeptide if it is expressed as a preprotein that participates in the secretion of the peptide or polypeptide; a promoter or enhancer is operably linked to a coding sequence if it affects the transcription of the sequence; or a ribosome binding site is operably linked to a coding sequence if it is positioned so as to facilitate translation. Generally, "operably linked" means that the DNA sequences being linked are contiguous and, in the case of a secretory leader, contiguous and in reading phase. However, enhancers do not have to be contiguous. Linking is accomplished by ligation at convenient restriction sites. If such sites do not exist, the synthetic oligonucleotide adaptors or linkers are used in accord with conventional practice.

The DNA to be introduced into the cells further will generally contain either a selectable marker gene or a reporter gene or both to facilitate identification and selection of transformed cells from the population of cells sought to be transformed. Alternatively, the selectable marker may be carried on a separate piece of DNA and used in a co-transformation procedure. Both selectable markers and reporter genes may be flanked with appropriate regulatory sequences to enable expression in the host cells. Useful selectable markers are well known in the art and include, for example, antibiotic and herbicide-resistance genes, such as *neo*, *hpt*, *dhfr*, *bar*, *aroA*, *dapA* and the like. See also, the genes listed on Table 1 of Lundquist et al. (U.S. Patent No. 5,848,956).

Reporter genes are used for identifying potentially transformed cells and for evaluating the functionality of regulatory sequences. Reporter genes which encode for easily assayable proteins are well known in the art. In general, a

reporter gene is a gene which is not present in or expressed by the recipient organism or tissue and which encodes a protein whose expression is manifested by some easily detectable property, e.g., enzymatic activity. Preferred genes include the chloramphenicol acetyl transferase gene (cat) from Tn9 of *E. coli*, the  
5 beta-glucuronidase gene (*gus*) of the *uidA* locus of *E. coli*, and the luciferase *luc* gene from firefly *Photinus pyralis*. Expression of the reporter gene is assayed at a suitable time after the DNA has been introduced into the recipient cells.

The general methods for constructing recombinant DNA which can transform target cells are well known to those skilled in the art, and the same  
10 compositions and methods of construction may be utilized to produce the DNA useful herein. For example, J. Sambrook et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory Press (2d ed., 1989), provides suitable methods of construction.

#### C. Transformation into Host Cells

15 The recombinant DNA can be readily introduced into the host cells, e.g., mammalian, bacterial, yeast or insect cells by transfection with an expression vector comprising DNA encoding TLR4 or its complement, by any procedure useful for the introduction into a particular cell, e.g., physical or biological methods, to yield a transformed cell having the recombinant DNA stably  
20 integrated into its genome, so that the DNA molecules, sequences, or segments, of the present invention are expressed by the host cell.

Physical methods to introduce a DNA into a host cell include calcium phosphate precipitation, lipofection, particle bombardment, microinjection, electroporation, and the like. Biological methods to introduce the DNA of  
25 interest into a host cell include the use of DNA and RNA viral vectors. The main advantage of physical methods is that they are not associated with pathological or oncogenic processes of viruses. However, they are less precise, often resulting in multiple copy insertions, random integration, disruption of foreign and endogenous gene sequences, and unpredictable expression. Viral  
30 vectors, useful to introduce genes to mammalian cells include, but are not limited to, poxvirus vectors, herpes simplex virus I vectors, adenovirus vectors, adeno-associated virus vectors, and the like.

As used herein, the term "cell line" or "host cell" is intended to refer to well-characterized homogenous, biologically pure populations of cells. These cells may be eukaryotic cells that are neoplastic or which have been "immortalized" *in vitro* by methods known in the art, as well as primary cells, or prokaryotic cells. The cell line or host cell is preferably of mammalian origin, but cell lines or host cells of non-mammalian origin may be employed, including plant, insect, yeast, fungal or bacterial sources. Generally, the DNA sequence is related to a DNA sequence which is resident in the genome of the host cell but is not expressed, or not highly expressed, or, alternatively, overexpressed.

"Transfected" or "transformed" is used herein to include any host cell or cell line, the genome of which has been altered or augmented by the presence of at least one DNA sequence, which DNA is also referred to in the art of genetic engineering as "heterologous DNA," "recombinant DNA," "exogenous DNA," "genetically engineered," "non-native," or "foreign DNA," wherein said DNA was isolated and introduced into the genome of the host cell or cell line by the process of genetic engineering. The host cells of the present invention are typically produced by transfection with a DNA sequence in a plasmid expression vector, a viral expression vector, or as an isolated linear DNA sequence.

To confirm the presence of the DNA sequence in the host cell, a variety of assays may be performed. Such assays include, for example, "molecular biological" assays well known to those of skill in the art, such as Southern and Northern blotting, RT-PCR and PCR; "biochemical" assays, such as detecting the presence or absence of a particular TLR4, e.g., by immunological means (ELISAs and Western blots) or by assays described herein.

To detect and quantitate RNA produced from introduced DNA segments, RT-PCR may be employed. In this application of PCR, it is first necessary to reverse transcribe RNA into DNA, using enzymes such as reverse transcriptase, and then through the use of conventional PCR techniques amplify the DNA. In most instances PCR techniques, while useful, will not demonstrate integrity of the RNA product. Further information about the nature of the RNA product may be obtained by Northern blotting. This technique demonstrates the presence of an RNA species and gives information about the integrity of that RNA. The presence or absence of an RNA species can also be determined using dot or slot

blot Northern hybridizations. These techniques are modifications of Northern blotting and only demonstrate the presence or absence of an RNA species.

While Southern blotting and PCR may be used to detect the DNA segment in question, they do not provide information as to whether the DNA segment is being expressed. Expression may be evaluated by specifically identifying the polypeptide products of the introduced DNA sequences or evaluating the phenotypic changes brought about by the expression of the introduced DNA segment in the host cell.

#### D. TLR4 polypeptides, variants, and derivatives thereof

- 10 The present isolated, purified TLR4 polypeptides, variants or derivatives thereof, can be synthesized *in vitro*, e.g., by the solid phase peptide synthetic method or by recombinant DNA approaches (see above). The solid phase peptide synthetic method is an established and widely used method, which is described in the following references: Stewart et al., Solid Phase Peptide  
15 Synthesis, W. H. Freeman Co., San Francisco (1969); Merrifield, J. Am. Chem. Soc., 85 2149 (1963); Meienhofer in "Hormonal Proteins and Peptides," ed.; C.H. Li, Vol. 2 (Academic Press, 1973), pp. 48-267; Bavaay and Merrifield, "The Peptides," eds. E. Gross and F. Meienhofer, Vol. 2 (Academic Press, 1980) pp. 3-285; and Clark-Lewis et al., Meth. Enzymol., 287, 233 (1997). These  
20 peptides can be further purified by fractionation on immunoaffinity or ion-exchange columns; ethanol precipitation; reverse phase HPLC; chromatography on silica or on an anion-exchange resin such as DEAE; chromatofocusing; SDS-PAGE; ammonium sulfate precipitation; gel filtration using, for example, Sephadex G-75; or ligand affinity chromatography.
- 25 Once isolated and characterized, derivatives, e.g., chemically derived derivatives, of a given TLR4 can be readily prepared. For example, amides of the TLR4 or TLR4 variants of the present invention may also be prepared by techniques well known in the art for converting a carboxylic acid group or precursor, to an amide. A preferred method for amide formation at the C-  
30 terminal carboxyl group is to cleave the peptide from a solid support with an appropriate amine, or to cleave in the presence of an alcohol, yielding an ester, followed by aminolysis with the desired amine.



Salts of carboxyl groups of a peptide or peptide variant of the invention may be prepared in the usual manner by contacting the peptide with one or more equivalents of a desired base such as, for example, a metallic hydroxide base, e.g., sodium hydroxide; a metal carbonate or bicarbonate base such as, for example, sodium carbonate or sodium bicarbonate; or an amine base such as, for example, triethylamine, triethanolamine, and the like.

N-acyl derivatives of an amino group of the TLR4 or variant thereof may be prepared by utilizing an N-acyl protected amino acid for the final condensation, or by acylating a protected or unprotected peptide. O-acyl derivatives may be prepared, for example, by acylation of a free hydroxy peptide or peptide resin. Either acylation may be carried out using standard acylating reagents such as acyl halides, anhydrides, acyl imidazoles, and the like. Both N- and O-acylation may be carried out together, if desired.

Formyl-methionine, pyroglutamine and trimethyl-alanine may be substituted at the N-terminal residue of the peptide or peptide variant. Other amino-terminal modifications include aminooxypentane modifications (see Simmons et al., *Science*, 276, 276 (1997)).

In addition, the amino acid sequence of TLR4 can be modified so as to result in a variant TLR4. The modification includes the substitution of at least one amino acid residue in the polypeptide for another amino acid residue, including substitutions which utilize the D rather than L form, as well as other well known amino acid analogs, e.g., unnatural amino acids such as  $\alpha$ ,  $\alpha$ -disubstituted amino acids, N-alkyl amino acids, lactic acid, and the like. These analogs include phosphoserine, phosphothreonine, phosphotyrosine, hydroxyproline, gamma-carboxyglutamate; hippuric acid, octahydroindole-2-carboxylic acid, statine, 1,2,3,4-tetrahydroisoquinoline-3-carboxylic acid, penicillamine, ornithine, citrulline,  $\alpha$ -methyl-alanine, para-benzoyl-phenylalanine, phenylglycine, propargylglycine, sarcosine,  $\epsilon$ -N,N,N-trimethyllysine,  $\epsilon$ -N-acetyllysine, N-acetylserine, N-formylmethionine, 3-methylhistidine, 5-hydroxylysine,  $\omega$ -N-methylarginine, and other similar amino acids and imino acids and tert-butylglycine.

Conservative amino acid substitutions include aspartic-glutamic as acidic amino acids; lysine/arginine/histidine as basic amino acids; leucine/isoleucine,

methionine/valine, alanine/valine as hydrophobic amino acids;  
serine/glycine/alanine/threonine as hydrophilic amino acids. Conservative  
amino acid substitution also includes groupings based on side chains. For  
example, a group of amino acids having aliphatic side chains is glycine, alanine,  
5 valine, leucine, and isoleucine; a group of amino acids having aliphatic-hydroxyl  
side chains is serine and threonine; a group of amino acids having amide-  
containing side chains is asparagine and glutamine; a group of amino acids  
having aromatic side chains is phenylalanine, tyrosine, and tryptophan; a group  
of amino acids having basic side chains is lysine, arginine, and histidine; and a  
10 group of amino acids having sulfur-containing side chains is cysteine and  
methionine. For example, it is reasonable to expect that replacement of a leucine  
with an isoleucine or valine, an aspartate with a glutamate, a threonine with a  
serine, or a similar replacement of an amino acid with a structurally related  
amino acid will not have a major effect on the properties of the resulting variant  
15 polypeptide. Whether an amino acid change results in a functional polypeptide  
can readily be determined by assaying the activity of the polypeptide variant.  
Such assays are described herein.

Conservative substitutions are shown in Figure 8 under the heading of  
exemplary substitutions. More preferred substitutions are under the heading of  
20 preferred substitutions. After the substitutions are introduced, the variants are  
screened for biological activity.

Amino acid substitutions falling within the scope of the invention, are, in  
general, accomplished by selecting substitutions that do not differ significantly  
in their effect on maintaining (a) the structure of the peptide backbone in the area  
25 of the substitution, (b) the charge or hydrophobicity of the molecule at the target  
site, or (c) the bulk of the side chain. Naturally occurring residues are divided  
into groups based on common side-chain properties:

- (1) hydrophobic: norleucine, met, ala, val, leu, ile;
- (2) neutral hydrophilic: cys, ser, thr;
- 30 (3) acidic: asp, glu;
- (4) basic: asn, gln, his, lys, arg;
- (5) residues that influence chain orientation: gly, pro; and
- (6) aromatic; trp, tyr, phe.

The invention also envisions polypeptide variants with non-conservative substitutions. Non-conservative substitutions entail exchanging a member of one of the classes described above for another.

5 Acid addition salts of the polypeptide or variant polypeptide or of amino residues of the polypeptide or variant polypeptide may be prepared by contacting the polypeptide or amine with one or more equivalents of the desired inorganic or organic acid, such as, for example, hydrochloric acid. Esters of carboxyl groups of the polypeptides may also be prepared by any of the usual methods known in the art.

#### 10 E. Methods of the Invention

The invention provides a method to diagnose individuals who are at a greater risk of deleterious consequences due to bacterial infection, e.g., the individuals may be more susceptible to infection by gram-negative bacteria, e.g., local gram-negative infection, more susceptible to sepsis induced by gram-  
15 negative bacteria, more susceptible to chronic airway disease, more susceptible to asthma, more susceptible to gall bladder disease, more susceptible to pyelonephritis, more susceptible to pneumonia, more susceptible to bronchitis, more susceptible to chronic obstructive pulmonary disease, more susceptible to arthritis, at higher risk for severe cystic fibrosis, and more susceptible to local  
20 and systemic inflammatory conditions such as systemic inflammatory response syndrome (SIRS), and acute respiratory distress syndrome (ARDS). The invention is also useful in the development of drugs that target the TLR4 gene product, e.g., increase or decrease the function of TLR4, especially the extracellular domain. These agents may thus be useful to prevent or ameliorate  
25 infection by gram-negative bacteria, prevent or ameliorate sepsis induced by gram-negative bacteria, prevent or ameliorate LPS-induced chronic airway disease in normal, cystic fibrosis and asthmatic populations, prevent or ameliorate arthritis, and prevent or ameliorate local and systemic inflammatory conditions such as SIRS and ARDS, particularly in individuals at risk for these  
30 indications or conditions.

The invention will be further described by the following non-limiting examples.

### Example 1

#### Methods

**Study Subjects.** The study population consisted of 83 healthy adult volunteers (31 men, 52 women) aged 18-50. Exclusion criteria included any history of allergies, tobacco use, cardiac or pulmonary disease. After written informed consent was obtained, all subjects were screened with spirometry, inhalation challenge with histamine, skin testing for common aeroallergens, chest x-ray, and electrocardiogram. All participants had normal screening studies (including histamine PC<sub>20</sub> > 32 mg/ml), were on no medications (except birth control), and had no significant acute or chronic cardiopulmonary disease or occupational exposures.

**Inhalation Challenge Protocol.** All subjects were exposed by inhalation challenge to buffered sterile saline (HBSS) followed by increasing concentrations of LPS. The solutions were delivered via a DeVilbiss 646 nebulizer powered by compressed air at 30 psi (DeVilbiss Co., Somerset, PA) and a Rosenthal dosimeter (Laboratory for Applied Immunology, Baltimore, MD). After the HBSS, subsequent inhalations delivered in increasing doses of LPS according to the following schedule: 0.5 µg, 1.0 µg, 2.0 µg, 3.0 µg, 5.0 µg, 10 µg, and 20 µg. Thus, the entire protocol delivered a total of 41.5 µg of LPS.

**Incremental LPS inhalation challenge.** The incremental LPS inhalation challenge was performed as follows. The % decline in FEV<sub>1</sub>/µg LPS was calculated following administration of the cumulative LPS dose that either resulted in at least a 20% decline in FEV<sub>1</sub> or the decline in FEV<sub>1</sub> following a cumulative inhaled dose of 41.5 µg of LPS. Subjects above the x-axis (solid bars in Figure 5) are homozygous for the wild type allele (WT/WT); subjects below the x-axis (open bars in Figure 5) are either heterozygous or homozygous (\*) for the missense Asp299Gly allele. The data is replotted in the inset after the values for the dose-response slope were log normalized. P values are presented for the comparison of the % decline in FEV<sub>1</sub>/µg LPS between subjects with the WT/WT genotype (N=73) and those with at least one Asp299Gly allele (N=10) using absolute values (P=0.037) and log normalized values (P=0.026). Since the distribution of the dose-response slope (% decline FEV<sub>1</sub>/cumulative dose of inhaled LPS) was highly skewed, the two-sample Monte-Carlo permutation test

based on 10,000 permutations was used to calculate P values (Fisher et al., 1993). To assess the allelic frequency of TLR4 sequence variants, a well-characterized Iowa population (Lidral et al., 1998) and the Centre d'Etude du Polymorphisme Humain (CEPH) population (NIH-CEPIT, 1992) were screened  
5 for specific sequence variants identified in the 83 study subjects.

Endotoxin. Solutions of endotoxin for inhalation were prepared according to a standard protocol using lyophilized *Escherichia coli* (serotype 0111:B4, Sigma Chemical Co., St. Louis, MO) LPS. These solutions of LPS were resuspended in sterile Hank's balanced salt solution (without calcium or magnesium) at a pH of  
10 7.0 and filter sterilized. All solutions used for inhalation were tested for sterility (bacteria and fungi) and LPS content (*Limulus ameobocyte* lysate assay, QCL-1000; Whittaker Bioproducts, Walkersville, MD) prior to separation into individual aliquots. These aliquots were stored immediately after preparation at -70°C until used.

15 Physiologic Measurements. A Spirotech (Atlanta, GA) S600 spirometer was used to assess pulmonary function; spirometry was performed using standards established by the American Thoracic Society. Subjects were positioned upright in a chair and were using noseclips. Baseline spirometry was recorded after inhalation of saline, and then 1, 10, 20, and 30 minutes following inhalation of  
20 each dose of LPS, and compared with the post-saline baseline spirometry. If the study subject's FEV<sub>1</sub> was greater than 80 % of the baseline measurement at the final assessment (30 minutes post-saline), the inhalation challenge was continued and the next does of LPS was administered. The challenge test was terminated when any of the following criteria had been met: 1) the subject did not wish to  
25 continue for any reason; 2) the subject's FEV<sub>1</sub> decreased 20 % or greater from baseline; or 3) a cumulative dose of 41.5 µg had been achieved. Of 84 subjects enrolled in the study, 1 subject withdrew prior to completion of the LPS inhalation challenge test, 52 subjects had at least a 20 % decline in the FEV<sub>1</sub> during the LPS inhalation challenge test, and 31 subjects inhaled a cumulative  
30 dose of 41.5 µg of LPS and did not decrease their FEV<sub>1</sub> by 20 %.

Assignment of Phenotype. Study subjects were categorized as either "responsive" or "hyporesponsive" to inhaled LPS. In the course of previous investigations, a large number of study subjects have been exposed to inhaled

- LPS (Jagiello et al., 1996; Deetz et al., 1997). In general, most healthy non-asthmatic study subjects develop airflow obstruction ( $FEV_1 \leq 80\%$  of the pre-exposure value) when challenged with approximately 40  $\mu\text{g}$  of LPS. Based on this and a standard approach to the definition of airway hyperreactivity ( $FEV_1$  decline of 20 % from pre-exposure values) (Chai et al., 1975), subjects were categorized as "LPS responsive" if they decreased their  $FEV_1$  by at least 20 % at any point during the LPS inhalation challenge, or "hypo-responsive" if they had < 20 % decline in their  $FEV_1$  after inhaling a cumulative dose of 41.5  $\mu\text{g}$  of LPS.
- Isolation of Genomic Clone. A human bacterial artificial chromosomal (BAC) library (Research Genetics; Huntsville, AL) was screened with two sets of primers (1F: 5'ATGGGGCATATCAGAGCCTA 3; SEQ ID NO:8, IR: 5'GTCCAATGGGGAAGTTCTCT 3; SEQ ID NO:9, 2F: 5'TCATTGTCCTGCAGAAGGTG 3; SEQ ID NO:10, and 2R: 5'CAGGGCTTTTCTGAGTCGTC 3; SEQ ID NO:11) derived from the human TLR4 gene (Genbank Accession Nos. U88880 and U93091). These sets of primers amplified a 160 bp and 140 bp product, respectively. PCR reactions were prepared by combining the following components: 1  $\mu\text{l}$  of PCR 10X buffer (100 mM Tris-HCl, pH 8.3, 500 mM KCl), 200  $\mu\text{M}$  each of dCTP, dGTP, dATP, and dTTP, 0.25  $\mu\text{M}$  of each primer, 0.2 U of *AmpliTaq* DNA polymerase (Perkin Elmer; Norwalk, CT) and 1  $\mu\text{l}$  of the library sample in a final volume of 10  $\mu\text{l}$ . Thermal cycling was performed with an initial denaturation at 94°C for 3 minutes followed by 35 cycles comprising 94°C, 55°C, and 72°C steps of 30 seconds each and a final extension of 72°C for 5 minutes. PCR products were separated by electrophoresis on 2% agarose gels, stained with ethidium bromide and visualized under UV light.
- Mutation Detection. Genomic DNA was isolated from whole blood obtained from the study subjects using a rapid salt isolation procedure (Laitinen et al., 1994). Overlapping primer sets were designed across the coding sequence such that products did not exceed 250 bp (Table 1). Primers were derived from flanking intronic sequences to include all splice sites. PCR reactions were prepared as described above except that 10-20 ng of genomic DNA was used as template. Amplification products were separated on non-denaturing, fan-cooled gels containing 5% acrylamide/bis (19:1), 0.5X TBE, and 2.5% glycerol for 3

hours at 20 W. A subset of PCR products were also run on MDE gels. The gels were subjected to silver staining and aberrant bands extracted from the gel, reamplified, and sequenced in both directions. To verify the sequence of the aberrant band, the same primers were used to amplify and sequence genomic

5 DNA from each subject. At least one individual without the aberrant band was also sequenced for comparison. Amplification products derived from the extracted DNA were purified using spin columns (Qiagen; Santa Clarita, CA) and DNA concentration determined by spectrophotometry. The DNA sequence was determined with a Model 377 automated DNA sequencer (Perkin Elmer;

10 Norwalk, CT).

Table 1. Primers for amplification of TLR4 coding region.

Exon 5' NT <sup>a</sup> FORWARD PRIMER (5'-3')		5' NT <sup>a</sup> REVERSE PRIMER (5'-3')		SIZE (bp)	
1	24	CTGCTCGGTCAAACGGTGAT <sup>1</sup>	+33	AGTCAGCTCCTCTGAACTTTC <sup>2</sup>	206
2	-106	CAGCAAGCACGATATTGGAT <sup>3,b</sup>	+18	CGTGCTCAGCATTTCTAACCT <sup>4,b</sup>	244
3	-53	GAGTTGGGAGACCATGCAGT <sup>5</sup>	+41	GAACACCTCACCTTGTGCAG <sup>6</sup>	262
4	-52	GGTTCTTATTCAGCAGAAAT <sup>7</sup>	483	CTTGATAGTCCAGAAAAGGCT <sup>8</sup>	172
4	496	GGTGGCTGTGGAGACAAATC <sup>9</sup>	691	CCGCAAGTCTGTGCAATAAA <sup>10</sup>	195
4	639	ACTTGGACCTTTCCAGCAAC <sup>11</sup>	798	GTACAGCTTATGAAGCCTAATTCT <sup>12</sup>	159
4	751	CTTTATCCAAACCAGGTGCA <sup>13</sup>	952	CAAAATTGCACAGGCCCTCTAGA <sup>14</sup>	201
4	882	TGGGAGAAATTTAGAAATGAAGGA <sup>15</sup>	1080	AAAGTCTTTTACCCTTTTCAATAGTCA <sup>16</sup>	198
4	1032	TTTCTTCATTTTCCCTGGTGA <sup>17</sup>	1161	AGAGATTTGAGTTTCAATGTGGG <sup>18</sup>	129
4	1129	TGGACAGTTTCCACACATTGA <sup>19</sup>	1309	CTTTAGGCTGGTTGTCCCAA <sup>20</sup>	180
4	1262	TTCAAAGGTTGCTGTTCTCAAA <sup>21</sup>	1426	TGAAAACCTCACTCATTTGTTTCAA <sup>22</sup>	164
4	1349	TCAAACCTTCTTGGGCTTAGAACA <sup>23</sup>	1526	CTTCGAGACTGGACAAGCCA <sup>24</sup>	177
4	1480	CAGAGTTGCTTTCAATGGCA <sup>25</sup>	1614	GAGAGTCCAGGAAGGTCAA <sup>26</sup>	134
4	1551	TCCAGGAAAACCTTCCCTCCA <sup>27</sup>	1746	ACCTGGAGGGAGTTCAGACA <sup>28</sup>	195
4	1702	TTCAATTGGATACGTTTCC <sup>29</sup>	1905	AAGAGCTGCCCTCTGGTCCCT <sup>30</sup>	203
4	1863	ACCAGAGTTTCCCTGCAATGG <sup>31</sup>	2032	TACAAGCACACTGAGGACCG <sup>32</sup>	169
4	1956	TGCCTGTGCTGAGTTTGAAT <sup>33</sup>	2100	TTTATGCAGCCAGCAAGAAG <sup>34</sup>	144
4	2013	CGGTCCCTCAGTGTGCTTGA <sup>35</sup>	2208	GGAGGCACCCCTTCTTCTAA <sup>36</sup>	195
4	2149	CCAGGATGAGGACTGGGTAA <sup>37</sup>	2338	GCGGCTCTGGATGAAGTGCT <sup>38</sup>	189
4	2290	AAGCCGAAAGGTGATTGTTG <sup>39</sup>	2442	CTGAGCAGGGTCTTCTCCAC <sup>40</sup>	152
4	2398*	TATCATCTTCAATTGTCTGCAGA <sup>41</sup>	2617	AGATGTTGCTTCTCGCCAA <sup>42</sup>	219
4	2406	TCATTGCTCTGCAGAAAGGTG <sup>43</sup>	2548	CAGGGCTTTTCTGAGTCGTC <sup>44</sup>	142
4	2529	GACGACTCAGAAAAGCCCTG <sup>45</sup>	2683	TGAACAAGTGTGGACCCAG <sup>46</sup>	154
4	2597	AATTGGCAGGAAGCAACATC <sup>47</sup>	2772	GATTAGCAGCCCTGCATATCT <sup>48</sup>	175



<sup>a</sup> relative to published cDNA sequence of Medzhitov et al. (1997), Genbank accession #U93091

<sup>b</sup> relative to published cDNA sequence of Rock et al. (1997), Genbank Accession #U88880, since exon 2 is absent in #U93091.

A "-" indicates primer is within intron on the 5' side of amplified exon; a "+" indicates primer is on the 3' side.

<sup>1</sup> SEQ ID NO:12	<sup>13</sup> SEQ ID NO:18	<sup>25</sup> SEQ ID NO:24	<sup>37</sup> SEQ ID NO:30
<sup>2</sup> SEQ ID NO:36	<sup>14</sup> SEQ ID NO:42	<sup>26</sup> SEQ ID NO:48	<sup>38</sup> SEQ ID NO:54
<sup>3</sup> SEQ ID NO:13	<sup>15</sup> SEQ ID NO:19	<sup>27</sup> SEQ ID NO:25	<sup>39</sup> SEQ ID NO:31
<sup>4</sup> SEQ ID NO:37	<sup>16</sup> SEQ ID NO:43	<sup>28</sup> SEQ ID NO:49	<sup>40</sup> SEQ ID NO:55
<sup>5</sup> SEQ ID NO:14	<sup>17</sup> SEQ ID NO:20	<sup>29</sup> SEQ ID NO:26	<sup>41</sup> SEQ ID NO:32
<sup>6</sup> SEQ ID NO:38	<sup>18</sup> SEQ ID NO:44	<sup>30</sup> SEQ ID NO:50	<sup>42</sup> SEQ ID NO:56
<sup>7</sup> SEQ ID NO:15	<sup>19</sup> SEQ ID NO:21	<sup>31</sup> SEQ ID NO:27	<sup>43</sup> SEQ ID NO:33
<sup>8</sup> SEQ ID NO:39	<sup>20</sup> SEQ ID NO:45	<sup>32</sup> SEQ ID NO:51	<sup>44</sup> SEQ ID NO:57
<sup>9</sup> SEQ ID NO:16	<sup>21</sup> SEQ ID NO:22	<sup>33</sup> SEQ ID NO:28	<sup>45</sup> SEQ ID NO:34
<sup>10</sup> SEQ ID NO:40	<sup>22</sup> SEQ ID NO:46	<sup>34</sup> SEQ ID NO:52	<sup>46</sup> SEQ ID NO:58
<sup>11</sup> SEQ ID NO:17	<sup>23</sup> SEQ ID NO:23	<sup>35</sup> SEQ ID NO:29	<sup>47</sup> SEQ ID NO:35
<sup>12</sup> SEQ ID NO:41	<sup>24</sup> SEQ ID NO:47	<sup>36</sup> SEQ ID NO:53	<sup>48</sup> SEQ ID NO:59

Multi-tissue cDNA Expression Screen. Human adult (Clontech #K1420-1; Palo Alto, CA) and fetal (Clontech #K1425-1; Palo Alto, CA) multi-tissue cDNA panels were screened by PCR using primers derived from exon 1 (forward; 5'GCTCACAGAAGCAGTGAGGA 3'; SEQ ID NO:60) and exon 4 (reverse; 5'TAGGCTCTGATATGCCCCAT3'; SEQ ID NO:61) of the human TLR4 gene. These PCR experiments were performed in 10 µl reactions composed of 0.025 U/µl of DNA polymerase (BioXACT; BioLine; Reno, NV), 1.0 µM of each primer, 200 µM of each dNTP, 1 µl of 10X buffer supplied by the manufacturer, and 1 µl of the cDNA sample. PCR conditions were: 95°C for 2 minutes; cycles of 0.5 minutes at 94°C, 0.5 minutes at 55°C, and 1 minute at 68°C; followed by a final extension step for 10 minutes at 68°C. The amplified products were separated on a 2% agarose gel, stained with ethidium bromide, and visualized under UV light.

Statistical Analysis. The statistical analysis was designed to determine whether specific mutations in the TLR4 gene were associated with the airway hyporesponsiveness to inhaled LPS. A one-tailed test of statistical significance was employed (Fleiss, 1986). In a 2 X 2 analysis, a Fisher's one-tailed exact test was used to determine whether specific mutations of the TLR4 gene occurred more frequently in study subjects who were considered LPS hyporesponsive compared to those with a normal airway response to inhaled LPS. In addition, because of the nonparametric distribution of dose-response slope (percent decline FEV<sub>1</sub>/cumulative dose of inhaled LPS), this outcome log was transformed and the Student's one-tailed T test (assuming unequal variances) was used to determine whether the dose-response slope was significantly less in subjects with a specific mutation of TLR4 compared to subjects with the common TLR4 allele.

Bioassays. Cells were maintained in F12 media supplemented with 10% fetal calf serum (Gibco, Rockville, MD), 2 mM L-glutamine, and 10,000 units penicillin/streptomycin, with 1 µg/ml of G418 as selective antibiotic. Cells were transfected with a mixture of 4 µg of DNA and 10 µl of Superfect (Qiagen, Valencia, CA) for a 35 mm<sup>2</sup> dish. The DNA mix consisted of 2 µg of DNA and 10 µg NFκB reporter plasmid encoding for the luciferase gene (Clontech, Palo Alto, CA) and 1 µg of each TLR4 (Medzhitov et al., 1997; Genbank #U93091)

expression plasmid (WT or Asp299Gly). If only one expression plasmid was used, empty vector pcDNA3.1<sup>+</sup> was added to keep the DNA concentration constant. Twenty-four hours later the cells were stimulated with 100 ng/ml of LPS for 6 hours. Total luciferase activity was measured using a commercially available method (Tropix, Bedford, MA). Briefly, after rising with PBS, cells were removed from filters by incubation with 120  $\mu$ l lysis buffer (25 mM Tris-phosphate, pH 7.8; 2 mM DTT; 2 mM 1, 2-diaminocyclohexane-N,N',N'-tetraacetic acid; 10% glycerol; and 1% Triton X-100) for 15 minutes. Light emission was quantified in a luminometer (Analytical Luminescence Laboratory, San Diego, CA).

Airway epithelial cells were obtained from trachea and bronchi of lungs removed for organ donation. Cells were genotyped for TLR4 and isolated by enzyme digestion as previously described (Zabner et al., 1996). Freshly isolated cells were seeded at a density of  $5 \times 10^5$  cells/cm<sup>2</sup> onto collagen-coated, 0.6 cm<sup>2</sup> diameter millicell polycarbonate filters (Millipore Corp., Bedford, MA). The cells were maintained at 37°C in a humidified atmosphere of 5% CO<sub>2</sub> and air. Twenty-four hours after plating, the mucosal media was removed and the cells were allowed to grow at the air-liquid interface. The culture media consisted of a 1:1 mix of DMEM/Ham's F12, 5% Ultraser G (Biopre SA, Cedex, France), 100 U/ml penicillin, 100  $\mu$ g/ml streptomycin, 1% nonessential amino acids, and 0.12 U/ml insulin. Epithelia were tested for transepithelial resistance, and for morphology by scanning electron microscopy. Fourteen days after seeding, the basal release of IL-1 $\alpha$  was measured in WT/WT (12 specimens from 4 individuals) and WT/Asp299Gly (24 specimens from 4 individuals) epithelia by collecting the basolateral conditioned media after 24 hours (Figure 6b). The epithelia were then exposed to 100 ng/ml of LPS on the apical side for 6 hours, and the basolateral media was collected after 24 hours. IL-1 $\alpha$  was measured using a commercially available ELISA (R&D; Minneapolis, MN).

To rescue the LPS hyporesponsive phenotype, heterozygote (WT/Asp299Gly) airway epithelia (Figure 6c) or homozygote (Asp299Gly/Asp299Gly) alveolar macrophages (Figure 6d) were transfected with a recombinant adenovirus vector expressing TLR4 (Rock et al. 1998; Genbank #U88880) that was prepared as described previously (Dandson et al.,

1994) by the University of Iowa Gene Transfer Vector Core at titers of about  $10^{10}$  infectious units (I.U.)/ml. Briefly TLR4 plasmid was blunt ended and cloned into the shuttle vector pAd5/CMV $\kappa$ -NpA using the EcoRV site. Fourteen days after seeding of the airway epithelia (20 specimens from 4 individuals), 50 MOI of the recombinant viruses (Ad/TLR4 and Ad/eGFP in phosphate buffered saline) were added to the basolateral surface of the epithelia for 30 minutes (Walters et al., 1999). After infection, the epithelia were incubated at 37°C for an additional 48 hours before the LPS stimulation assay. To assay for basal LPS response, the media was changed with fresh 500  $\mu$ l of Ultrosor G, and collected after 24 hours to measure the basal IL-1 $\alpha$  secretion to the basolateral side. After collecting the basal specimen, the epithelia were exposed to 100 ng/ml of LPS on the apical side for 6 hours, and the media was collected after 24 hours. To assay for gene transfer efficiency, the epithelia was associated with 0.05% trypsin and 0.53 mM EDTA. Fluorescence from 50,000 individual cells was analyzed using fluorescence-activated cell analysis (FACScan, Lysys II software, Becton Dickinson, San Jose, CA). The percentage of GFP positive cells ranged between 52% and 76%. Human alveolar macrophages were collected by BAL from a homozygote (Asp299Gly/Asp299Gly) study subject as previously described (Deetz et al., 1967). Macrophages were seeded onto a 96 well plate at a density of  $10^5$  cells per well. The cells were infected after 4 hours with Ad/TLR4 in a CaPi coprecipitate (Iasbinder et al., 1998) at an MOI of 50, and a  $\text{Ca}^{++}$  concentration of 16.8 mM. Sixteen hours after infection, the cells were exposed to LPS in 1% serum for 6 hours, and the media was collected. TNF- $\alpha$  was measured using a commercially available ELISA (R&D; Minneapolis, MN).

## 25 Results

Genomic Structure of the TLR4 Gene. To determine the genomic structure of the TLR4 gene, a bacterial artificial chromosome (BAC) library was screened by PCR with primers derived from the 5' and 3' ends of the cDNA sequence. Human BAC clone 439F3 was identified with both sets of primers and sequenced to identify exon/intro splice sites using the two published cDNA sequences (Medzhitov et al., 1997; Rock et al., 1998). The TLR4 cDNA published by Medzhitov et al. (1997) (Genbank Accession No. U93091) was missing a 120 bp sequence beginning at nucleotide 136 of the other published

cDNA (Rock et al., 1998) (Genbank Accession No. U88880). Assuming this missing sequence represented an alternatively spliced exon, DNA sequencing primers were designed to amplify across the predicted splice junctions. When the genomic sequence (Figure 9) was compared to the cDNA sequences, the exon/intron junctions were revealed (Figure 1). The human TLR4 gene is organized into 4 exons that span about 11 kb of genomic DNA. As predicted, the 120 bp sequence missing in the cDNA from Medzhitov et al. (1997) corresponds to exon 2.

To determine whether other exons may be revealed by additional splice forms, a panel of cDNAs from adult (Figure 2a) and fetal (Figure 2b) tissues were examined with PCR primers designed to amplify all nucleotides between exons 1 and 4. In all adult and fetal tissues, three products were amplified (453, 333, and 167 bp), although their relative amounts varied. In addition to the 3' end of exon 1 and the 5' end of exon 4, the 453 bp product contained both exons 2 and 3, the 333 bp product was missing exon 2, and the 167 bp product was missing both exons 2 and 3 (Figure 2c). The 453 and 333 bp products are identical to the published cDNA sequences (Medzhitov et al., 1997; Rock et al., 1998), and confirm that no other exons are present for the human TLR4 gene. The 167 bp product depicts alternative splicing of exons 1 and 4, and represents a novel TLR4 product. The nucleotide position of alleles are relative to the TLR4 cDNA sequence published by Medzhitov et al. (1997).

**Mutation Analysis.** SSCV was employed to detect sequence variants across the entire coding region of TLR4 gene in the 83 unrelated probands who completed the incremental LPS inhalation challenge test. When band shifts were detected on SSCV analysis, the bands were sequenced to identify and confirm the sequence variants. The SSCV and sequence analysis were performed blinded to the LPS response phenotype of the study subjects. A band variant was detected by SSCV in 10 (12 %) of the 83 subjects, and direct sequencing detected an A to G substitution at nucleotide 896 of the published TLR4 cDNA (Figure 3a) (Genbank Accession No. U88880). To confirm these findings, the 83 unrelated probands were sequenced in the forward and reverse directions with primers designed to amplify the 896 nucleotide; the same 10 individuals were found to have the A896G substitution and the remaining individuals were confirmed to

have the common TLR4 allele. Importantly, one of the 10 individuals with the A896G substitution was homozygous for this mutation and the remaining 9 had a single mutant allele. The allelic frequency of the A896G substitution was 6.6 % in the study population, 7.9 % in a normal control population from Iowa (Lidral et al., 1998), and 3.3 % in the parental chromosomes of the CEPH population (NIH-CEPH, 1992).

The A896G substitution results in replacement of a conserved aspartic acid (A) residue with glycine (G) at amino acid 299 (Figure 4). This missense mutation (Asp299Gly) is in the fourth exon of TLR4 and is present in the extracellular domain of this receptor. The region surrounding amino acid 299 appears to be in an alpha helical conformation (Gibrat et al., 1987). Replacement of the conserved aspartic acid with glycine at position 299 causes disruption of the alpha helical protein structure resulting in the formation of an extended beta strand (Gibrat et al., 1987).

Two other SSCV variants were identified in a single proband; a T to C change at nucleotide 479 (Figure 3b) and a deletion of a thymine nucleotide at position -11 in the third intron. The T479C variant did not alter the amino acid composition and is therefore considered a silent mutation. The T-11 variant did not alter the acceptor splice site, so the significance of this deletion is unknown. The subject with these two sequence variants was responsive to inhaled LPS.

An additional missense mutation was identified within exon 4 of the TLR4 gene. This mutation results in the replacement of a threonine with an isoleucine residue at amino acid 399, which is on the 3' side of the previously identified Asp299Gly mutation. Both mutations are present within the extracellular domain of the TLR4 protein. When the genotypes were compared with the phenotypes associated with endotoxin responsiveness, one individual who carried the Asp299Gly mutation did not carry the Thr399Ile mutation. Likewise, one individual who had the Thr399Ile mutation did not have the Asp299Gly mutation. Nine of the 83 volunteers carry both mutations. There was a strong correlation between both of the TLR4 mutations and endotoxin hyporesponsiveness.

Phenotype/Genotype Analysis. Of the 83 unrelated study subjects who completed the LPS inhalation challenge test, 52 (63 %) were responsive to

inhaled LPS and 31 (37 %) were hyporesponsive to inhaled LPS. When the genotypes of these individuals were examined, the Asp299Gly sequence variant occurred in 3 LPS responsive (5.8 %) and 7 LPS hyporesponsive (22.6 %) study subjects ( $P=0.029$ ). Among the subjects with the common TLR4 allele ( $N = 73$ ), the dose-response slope (percent decline  $FEV_1$ /cumulative dose of inhaled LPS) averaged 1.86 % decline in  $FEV_1/\mu g$  inhaled LPS (range 0.01 %-19.78 %), while the dose-response slope for the subjects with the Asp299Gly allele ( $N = 10$ ) was significantly less ( $P=0.037$ ), averaging 0.59 % decline in  $FEV_1/\mu g$  inhaled LPS (range 0.00 %-1.59 %) (Figure 5). This group difference in the dose-response slope also exists on the log scale ( $P=0.026$ ) where  $\log(1 + \Delta FEV_1)$  was used since one individual has a  $\Delta FEV_1$  of 0.0. The subject who was homozygous for the Asp299Gly allele was hyporesponsive to inhaled LPS with a 0.28 % decline in  $FEV_1/\mu g$  of inhaled LPS. This homozygote Asp299Gly subject is one of a monozygote twin pair; her twin sister was subsequently phenotyped and was also found to be hyporesponsive to inhaled LPS with a 0.34% decline in  $FEV_1/\mu g$  of inhaled LPS.

The biological significance of the Asp299Gly mutation was evaluated in several ways. First, transfection of CHO cells with either the WT or the mutant TLR4 gene demonstrated that the mutant allele does not respond normally to LPS stimulation (Figure 6a). Second, airway epithelia obtained from heterozygote (WT/Asp299Gly) individuals do not respond to LPS stimulation (Figure 6b). Third, the wild-type allele of TLR4 clearly restored LPS responsiveness in either airway epithelial cells (Figure 6c) or alveolar macrophages (Figure 6d) obtained from individuals with the TLR4 mutation.

## Discussion

The results described herein provide the first direct evidence to indicate that a sequence polymorphism in the TLR4 gene is associated with a hyporesponsive LPS phenotype in humans that interrupts LPS signaling. This conclusion is supported by the following findings: 1) unrelated subjects with the Asp299Gly substitution were significantly less responsive to inhaled LPS than those homozygous for the common TLR4 allele; 2) an individual who was homozygous for the Asp299Gly substitution was hyporesponsive to inhaled LPS; 3) a monozygous twin pair, homozygous for the Asp299Gly substitution,

demonstrated a very similar response to inhaled LPS; and 4) *in vitro* studies demonstrate that mutant TLR4 does not respond to LPS stimulation. This conclusion is indirectly supported by the finding in C3H/HeJ mice where a point mutation in intracellular domain of *Tlr4* is also associated with LPS

- 5 hyporesponsiveness (Poltorak et al., 1998; Qureshi et al., 1999). The identified Asp299Gly missense mutation is in the fourth exon of the TLR4 gene and, unlike the C3H/HeJ mutation, is located in the extracellular domain of this receptor. The findings described herein suggest that a specific region in the extracellular domain of TLR4 may play an important role in receptor function
- 10 and regulation of the innate immune response to LPS in humans.

This discovery may have important ramifications for a broad spectrum of human diseases. First, the Asp299Gly TLR4 substitution is a relatively common mutation, potentially affecting the responsiveness of a substantial portion of the population to LPS. The allelic frequency of the Asp299Gly TLR4 substitution in

15 the three populations screened was between 3.3 % and 7.9 %, and homozygous individuals were identified in each of the populations. Second, the results described herein provide further support for the role of TLR4 in LPS signaling. A fundamental understanding of LPS signaling will undoubtedly alter approaches to gram-negative sepsis, as well as other diseases thought to be

20 mediated by endotoxin, such as the systemic inflammatory response syndrome (Wang et al., 1995), acute respiratory distress syndrome (Brighan et al., 1986), and asthma or other forms of airway disease caused (Rylander et al., 1989; Schwartz et al., 1995) or exacerbated (Schwartz et al., 1995; Michel et al., 1996) by endotoxin. Identification of the essential components of LPS signaling also

25 provide new therapeutic targets for endotoxin mediated conditions. Third, the specific Asp299Gly TLR4 substitution may provide a simple screening mechanism to risk stratify a population. Identification of this mutation may lead to a better appreciation of the role of LPS responsiveness in a broad range of acquired and genetic disorders. For example, mutations of the TLR4 gene may

30 explain why a minority of patients with gram-negative sepsis develop ARDS (Hudson et al., 1995) or why patients with cystic fibrosis and the identical CFTR mutation have different courses of disease (Veeze et al., 1994). Finally, it must be acknowledged that normal responsiveness to endotoxin is an important



component of innate immunity, and the Asp299Gly TLR4 substitution may prove to be associated with negative clinical outcomes. In fact, the C3H/HeJ mouse is more susceptible to *Salmonella typhimurium* (O'Brien et al., 1980), despite its resistance to LPS (Sultzcr et al., 1968). Although individuals with the

5 Asp299Gly TLR4 substitution may be more resistant to localized forms of endotoxin-induced inflammation, these individuals may prove to be more susceptible to a systemic inflammatory response initiated or exacerbated by endotoxin.

The findings described herein demonstrate that a specific region in the

10 extracellular domain of TLR4 plays an important role in receptor function and regulation of the innate immune response to LPS in humans. The sequence variants reported thus far for mammalian, i.e., murine, TLR4 are located in the cytoplasmic domain (Poltorak et al., 1998; Qureshi et al., 1999). While it is logical to hypothesize that mutations in the cytoplasmic domain of TLR4 disrupt

15 the signaling pathway leading to activation of NF- $\kappa$ B and AP-1 (Medhitov et al., 1998), the potential mechanisms that are altered by mutations in extracellular domain of TLR4 are less obvious.

TLR4 is a transmembrane protein and it is therefore not surprising to find a putative signal peptide at the N-terminus, presumably responsible for proper

20 trafficking to the cell membrane (Medzhitov et al., 1997). This finding raises the possibility that sequence variants in the extracellular domain of TLR4 can disrupt trafficking of this receptor to the cell membrane and enhance proteolysis. There are several examples in the literature which show that the presence of a single point mutation at a conserved amino acid can disrupt protein folding and

25 specifically affect protein trafficking (Jackson et al., 1998; DeFranco et al., 1998). Mutations of this kind could give rise to a range of phenotypes from almost normal to completely unresponsive, depending on the severity of the trafficking defect. Another possible mechanism that may explain the association between a mutation in the extracellular domain of TLR4 and LPS responsiveness

30 is impairment of ligand binding to the extracellular domain. The extracellular domains of the toll proteins are clearly involved in the cellular response to LPS (Medzhitor et al., 1997; Yang et al., 1998). Yet, so far no convincing evidence has been found that the toll proteins are directly interacting with LPS (Wright,

1999). The discovery of an association between a mutation in the extracellular domain of TLR4 and a discernable LPS phenotype in humans should aid in clarifying whether the extracellular domain of this receptor is important for proper membrane localization of TLR4 and if LPS directly interacts with TLR4.

- 5 A final possibility is that the missense mutation of TLR4 affects its interaction with either an extracellular ligand or an intracellular protein (Wright, 1999). The extracellular domain of TLR4, and specifically the Asp299Gly amino acid change may provide a key therapeutic target to modulate LPS signaling.

- TLR4 may be one of several factors that may regulate the airway
- 10 response to inhaled LPS. Among the subjects with the Asp299Gly substitution, 7 were hyporesponsive to LPS and 3 developed airflow obstruction during the LPS inhalation challenge test. None of the subjects with this mutation were in the most sensitive quartile of the LPS responders. Although asthmatic patients were included because of their increased airway responsiveness to inhaled LPS
- 15 (Michel et al., 1989), it remains possible that the 3 subjects with the Asp299Gly substitution who were responsive to inhaled LPS had some type of airway inflammation (e.g., early viral illness) that enhanced their response to inhaled LPS. Moreover, since 22.6 % of the study subjects who were hyporesponsive to LPS had at least one copy of the Asp299Gly TLR4 allele, other genes (or
- 20 possibly acquired conditions) may prove to play a role in modulating the biological response to LPS. For instance, TNF- $\alpha$  is one of the primary cytokines that mediates the toxic effects of LPS (Beutler et al., 1985). A polymorphism at -308 in the TNF- $\alpha$  promoter region results in higher constitutive and inducible levels of TNF- $\alpha$  (Wilson et al., 1997), and this polymorphism has been shown to
- 25 result in a worse outcome in children with either meningococcal disease (Nadel et al., 1996) or cerebral malaria (McGuire et al., 1994). Likewise, allelic variants of TLR4 receptors and their ligands appear to delay the progression of disease in patients with HIV infection (Mummidi et al., 1995). Thus, TLR4 appears to represent only one of a number of genetic variants that may modulate the
- 30 pathophysiologic response to LPS.

## Example 2

### Methods

- To genotype patients for TLR4, a PCR based RFLP assay was employed. For example, to detect point mutations in codon 299 such as those that result in an amino acid substitutions at amino acid 299 (Asp (GAT) to Gly (GGT) change) and/or in codon 399 (Thr (ACC) to Ile (ATC) change), a primer spanning that codon and a second primer 3' or 5' to that primer are prepared. Primers useful to detect a nucleotide change at codon 299 are: 299Forward primer (5' GATTAGCATACTTAGACTACTACCTCCATG 3'; SEQ ID NO:66) and 299Reverse primer (5' GATCAACTTCTGAAAAAGCATTCCCAC; SEQ ID NO:67). The underlined base in the forward primer fortuitously creates a NcoI (CCATGG) restriction site in the context of a mutation which is specific for carriers of the 299 mutation Asp to Gly. Primers useful to detect a nucleotide change at codon 399 include: 399Forward primer (5' GGTTGCTGTTCTCAAAGTGATTTTGGG AGA A; SEQ ID NO:68) and 399Reverse Primer (5' ACCTGAAGACTGGAGAGTGAGTTAAATGCT; SEQ ID NO:69). The underlined base in the 399 Forward primer fortuitously creates a HinfI restriction site (GANTC) in the context of a mutation which is specific for carriers of the 399 mutation Thr to Ile. The reverse primers were chosen at random based solely on such parameters as annealing temperature, expected product size and the like. Additional residues or fewer residues at the 5' end of the primers having the nucleotide change will not affect the outcome as long as the amplification conditions and reverse primers are chosen accordingly.
- A MJ Tetrad PTC-225 Thermo cycler was used with both primer sets. For the 299 assay, primers at 10-100 pmole were mixed with 20 ng of genomic DNA, and Perkin Elmer Taq polymerase, deoxyribonucleotides and buffer according to the manufacturer's directions. Cycling conditions included an initial denaturation of 95°C, for 3 to 4 minutes, then 30 cycles of denaturation at 95°C for 30 seconds; annealing at 55°C for 30 seconds; and extension at 72°C for 30 seconds.

For the 399 assay, a Clontech Advantage amplification kit was used according to the manufacturer's directions. The cycling conditions included an

initial denaturation at 95°C for 3 to 4 minutes, then 30 cycles of denaturation at 95°C for 30 seconds; annealing at 55°C for 30 seconds; and extension at 72°C for 30 seconds.

### Results

5        Septic shock remains a significant health concern world-wide and despite progress in understanding the physiologic and molecular basis of septic shock, the high mortality rate of septic shock patients remains unchanged. A common polymorphism in TLR4 was identified that is associated with hyporesponsiveness to inhaled endotoxin or lipopolysaccharide (LPS) in humans  
10 (Example 1). Since TLR4 is a major receptor for LPS in mammals and Gram-negative bacteria are the prevalent pathogen associated with septic shock, these specific TLR4 alleles may be associated with a predisposition to or a more severe disease outcome for septic shock patients. Ninety-one septic shock patients as well as 83 healthy blood donor controls were genotyped for the  
15 presence of the TLR4 Asp299Gly and TLR4 Thr399Ile mutations. The TLR4 Asp299Gly allele was found exclusively in septic shock patients ( $P = 0.05$ ). Furthermore, septic shock patients with the TLR4 Asp299Gly allele had a significantly higher prevalence of Gram-negative infections. These findings suggest that the TLR4 Asp299Gly allele predisposes people to develop septic  
20 shock with Gram-negative organisms.

      In another study in a German population, the following observations were made: patients with TLR4 mutation(s) tended to have a longer ICU stay; patients with TLR4 mutation(s) were more prone to sepsis; and sepsis in patients with TLR4 mutation(s) was more severe than in patients that were wild-  
25 type for TLR4.

      In a third study in a Finnish population, it was observed that mothers of pre-term infants and pre-term infants had a higher frequency of TLR4 mutation(s) than term infants. Also, infants with respiratory distress syndrome (RDS), particularly those born very prematurely, tended to have a higher  
30 frequency of mutation(s) than premature infants without respiratory distress syndrome.

Thus, generally, the 299 mutation is slightly more prevalent than the 399 mutation, and in 90% of the cases, both mutations were observed. When only one mutation was found, it was more frequently the 299 mutation.

Thus, the presence of TLR4 mutations is associated with predisposition  
 5 to septic shock, severity of sepsis, pre-term delivery, and RDS in pre-term infants.

#### References

- Beutler et al., Science, 229:869 (1985).
- 10 Bonner et al., Am. J. Respir. Cell Mol. Biol., 19:672 (1998).
- Brigham et al., Am. Rev. Respir. Dis. 133:913 (1986).
- Chai et al., J. Allergy Clin. Immunol., 56:323 (1975).
- Chow et al., J. Biol. Chem., 274:10689 (1999).
- Coutinho et al., Immunogenetics, 7:17 (1978).
- 15 Davidson et al., Exp. Neurol., 125:258 (1994).
- Deetz et al., Am. J. Respir. Crit. Care Med., 155:254 (1997).
- DeFranco et al., Prog. Clin. Biol. Res., 397:119 (1998).
- De Franco et al., J. Steroid Biochem. Mol. Biol., 65:51 (1998).
- Fasbender et al., J. Clin. Invest., 102:184 (1998).
- 20 Favorite et al., J. Clin. Invest., 21:589 (1942).
- Fisher et al., Wiley, New York (1993).
- Fleiss, J.C. The design and analysis of clinical experiments. New York:Wiley, 1986.
- Gibrat et al., J. Mol. Biol., 198:425 (1987).
- 25 Hoshino et al., J. Immunol., 162:3749 (1999).
- Hudson et al., Am. J. Respir. Crit. Care Med., 151:293 (1995).
- Jackson et al., Thromb. Haemost., 80:42 (1998).
- Jagiello et al., Chest, 110:263 (1996).
- Kuhns et al., J. Immunol., 158:3959 (1997).
- 30 Laitinen et al., Biotechniques, 17:316 (1994).
- Lidral et al., Am. J. Hum. Genet., 63:557 (1998).
- McGuire et al., Nature, 371:508 (1994).
- Medzhitov et al., Cell, 91:295 (1997).

- Medzhitov et al., Nature, 388:394 (1997).
- Medzhitov et al., Mol. Cell., 2:253 (1998).
- Michel et al., J. Appl. Physiol., 66:1059 (1989).
- Michel et al., Rev. Respir. Dis., 146:352 (1992).
- 5 Michel et al., Am. J. Respir. Crit. Care Med., 154:1641 (1996).
- Mummidi et al., Nat. Med., 4:786 (1998).
- Nadel et al., J. Infect. Dis., 174:878 (1996).
- NIH-CEPH (Collaborative Mapping Group), Science 258:67 (1992).
- O'Brien et al., Nature, 287:440 (1980).
- 10 Poltorak et al., Blood Cells Mol. Dis., 24:340 (1998a).
- Poltorak et al., Science, 282:2085 (1998).
- Qureshi et al., J. Exp. Med., 189:615 (1999).
- Rock et al., Proc. Natl. Acad. Sci. USA, 95:588 (1998).
- Rylander et al., Am. Rev. Respir. Dis., 140:981 (1989).
- 15 Santamaria et al., J. Immunol., 143:913 (1989).
- Schwartz et al., Am. J. Respir. Crit. Care Med., 151:47 (1995a).
- Schwartz et al., Am. J. Respir. Crit. Care Med., 152:603 (1995b).
- Sigmund et al., Hypertension, 22:599 (1993).
- Sultzter, Nature, 219:1253 (1968).
- 20 Vogel et al., J. Immunol., 122:619 (1979).
- Veeze et al., J. Clin. Invest., 93:461 (1994).
- Walters et al., J. Biol. Chem., 274:10219 (1999).
- Wang et al., Clin. Nucl. Med., 20:494 (1995).
- Wilson et al., Proc. Natl. Acad. Sci. USA, 94:3195 (1997).
- 25 Wright et al., Science, 249:1431 (1990).
- Wright et al., J. Exp. Med., 189:605 (1999).
- Yang et al., Nature, 395:284 (1998).
- Zabner et al., J. Virol., 70:6694 (1996).

30. All publications, patents and patent applications are incorporated herein by reference. While in the foregoing specification, this invention has been described in relation to certain preferred embodiments thereof, and many details have been set forth for purposes of illustration, it will be apparent to those skilled

in the art that the invention is susceptible to additional embodiments and that certain of the details herein may be varied considerably without departing from the basic principles of the invention.

**WHAT IS CLAIMED IS:**

1. An isolated and purified DNA molecule comprising a human genomic DNA segment encoding TLR4, a biologically active fragment or variant thereof.
2. The DNA molecule of claim 1 wherein the DNA segment encodes a variant TLR4.
3. The DNA molecule of claim 2 wherein the variant TLR4 has at least one amino acid substitution in the extracellular domain of TLR4.
4. The DNA molecule of claim 3 wherein the variant TLR4 comprises a glycine residue at position 299 or an isoleucine residue at position 399.
5. An isolated and purified nucleic acid molecule comprising a human nucleic acid segment encoding a variant TLR4, or a biologically active fragment thereof.
6. The nucleic acid molecule of claim 5 wherein the variant has at least one amino acid substitution in the extracellular domain of TLR4.
7. The nucleic acid molecule of claim 6 wherein the variant comprises a glycine residue at position 299 or an isoleucine residue at position 399.
8. An expression cassette comprising a promoter operably linked to a human DNA segment encoding a variant TLR4, or a biologically active fragment thereof.
9. The expression cassette of claim 8 wherein the variant TLR4 has at least one amino acid substitution in the extracellular domain of TLR4.



10. The expression cassette of claim 8 wherein the variant TLR4 comprises a glycine residue at position 299 or an isoleucine residue at position 399.
11. A host cell, the genome of which is augmented by the DNA molecule of claim 1.
12. A host cell, the genome of which is augmented by the nucleic acid molecule of claim 5.
13. A method to prepare a variant TLR4, comprising: culturing a host cell transformed with the nucleic acid molecule of claim 5 operably linked to a promoter, so that said host cell expresses the variant TLR4.
14. A method to prepare a variant TLR4, comprising: culturing a host cell transformed with the DNA molecule of claim 1 operably linked to a promoter, so that said host cell expresses the variant TLR4.
15. The method of claim 13 or 14 further comprising isolating TLR4 from the host cell.
16. A method of identifying a human at risk of, or having, an indication associated with altered innate immunity, comprising:
  - a) contacting an amount of DNA obtained from a human physiological sample with an amount of at least one TLR4-specific oligonucleotide under conditions effective to amplify the DNA so as to yield amplified DNA; and
  - b) detecting or determining whether the amplified DNA encodes at least a portion of a variant TLR4.
17. The method of claim 16 wherein the nucleic acid sample comprises genomic DNA.
18. The method of claim 16 wherein the nucleic acid sample is cDNA.

19. The method of claim 16 wherein the amplified DNA is subjected to electrophoresis in step b).
20. Isolated, variant human TLR4 polypeptide, or a biologically active fragment thereof.
21. The isolated polypeptide of claim 20 wherein the variant TLR4 has at least one amino acid substitution in the extracellular domain of TLR4.
22. The isolated polypeptide of claim 21 wherein the variant TLR4 comprises a glycine residue at position 299 or an isoleucine residue at position 399.
23. A method to identify an agent that alters TLR4 activity, comprising:
  - a) contacting a mammalian cell which expresses TLR4 with the agent;and
  - b) detecting or determining whether the agent alters TLR4 activity.
24. The method of claim 23 wherein the mammalian cell is transformed with a recombinant nucleic acid molecule encoding human TLR4.
25. An agent identified by the method of claim 24.
26. A method to alter TLR4 activity, comprising: contacting a mammalian cell with an agent effective to alter TLR4 activity.
27. The method of claim 26 wherein the mammalian cell is a human cell.
28. A method to detect a polymorphism in a human TLR4 gene, comprising: detecting or determining whether a nucleic acid sample obtained from a human comprises nucleic acid encoding variant TLR4.

29. A transgenic mouse whose cells comprise a chimeric DNA sequence, wherein the chimeric DNA sequence comprises a DNA segment encoding human TLR4, wherein the chimeric DNA sequence is integrated into the genome of the cells of the mouse, and wherein the DNA segment is expressed in the transgenic mouse so as to result in the transgenic mouse exhibiting altered innate immunity relative to a corresponding nontransgenic mouse.
30. The transgenic mouse of claim 29 wherein the DNA segment encodes a variant human TLR4.
31. The method of claim 16 wherein the TLR4-specific oligonucleotide comprises at least two nucleotide substitutions relative to the wild-type TLR4 nucleic acid sequence, which substitutions result in a restriction site that is not present in the wild-type TLR4 nucleic acid sequence and which restriction site is indicative of DNA that encodes a variant TLR4.
32. The method of claim 31 wherein the DNA encodes a variant TLR4 having an amino acid substitution at residue 299.
33. The method of claim 31 wherein the oligonucleotide comprises SEQ ID NO:66.
34. The method of claim 31 wherein the DNA encodes a variant TLR4 having an amino acid substitution at residue 399.
35. The method of claim 31 wherein the oligonucleotide comprises SEQ ID NO:68.

## SEQUENCE LISTING

<110> University of Iowa Research Foundation

5 Schwartz, David A.

Schutte, Brian C.

<120> Variant TLR4 nucleic acid and uses thereof

10<130> 875.010W01

<150> US 09/329,515

<151> 1999-06-10

15<160> 72

<170> FastSEQ for Windows Version 4.0

<210> 1

20<211> 28

<212> DNA

<213> Homo sapiens

<400> 1

25gcgtggaggt atgtggctgg agtcagct

28

<210> 2

<211> 28

<212> DNA

30<213> Homo sapiens

<400> 2

tcacggaggt tagaatgctg agcacgta

28

35<210> 3

<211> 28

<212> DNA

<213> Homo sapiens

40<400> 3

ttatccaggt aatgaatcca cttttaca

28

2

&lt;210&gt; 4

&lt;211&gt; 21

&lt;212&gt; PRT

&lt;213&gt; Homo sapiens

5

&lt;400&gt; 4

Leu Ala Tyr Leu Asp Tyr Tyr Leu Asp Asp Ile Ile Asp Leu Phe Asn

1

5

10

15

Cys Leu Thr Asn Val

10

20

&lt;210&gt; 5

&lt;211&gt; 20

&lt;212&gt; PRT

15&lt;213&gt; Mus musculus

&lt;400&gt; 5

Leu Thr Tyr Thr Asn Asp Phe Ser Asp Asp Ile Val Lys Phe His Cys

1

5

10

15

20Leu Ala Asn Val

20

&lt;210&gt; 6

&lt;211&gt; 20

25&lt;212&gt; PRT

&lt;213&gt; Rattus norvegicus

&lt;400&gt; 6

Leu Thr Tyr Ile Asn His Phe Ser Asp Asp Ile Tyr Asn Leu Asn Cys

30 1

5

10

15

Leu Ala Asn Ile

20

&lt;210&gt; 7

35&lt;211&gt; 20

&lt;212&gt; PRT

&lt;213&gt; Cricetulus griseus

&lt;400&gt; 7

40Phe Thr Tyr Ala Asn Glu Phe Ser Glu Asp Ile Thr Asp Phe Asp Cys

3

1

5

10

15

Leu Ala Asn Val

20

5&lt;210&gt; 8

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

10&lt;400&gt; 8

atggggcata tcagagccta

20

&lt;210&gt; 9

&lt;211&gt; 20

15&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 9

gtccaatggg gaagttctct

20

20

&lt;210&gt; 10

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25

&lt;400&gt; 10

tcattgtcct gcagaagggtg

20

&lt;210&gt; 11

30&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 11

35cagggctttt ctgagtcgtc

20

&lt;210&gt; 12

&lt;211&gt; 20

&lt;212&gt; DNA

40&lt;213&gt; Homo sapiens

&lt;400&gt; 12

ctgctcggtc aaacggtgat

20

&lt;210&gt; 13

5&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 13

10cagcaagcac gatattggat

20

&lt;210&gt; 14

&lt;211&gt; 20

&lt;212&gt; DNA

15&lt;213&gt; Homo sapiens

&lt;400&gt; 14

gagttgggag accatgcagt

20

20&lt;210&gt; 15

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25&lt;400&gt; 15

ggttcttatt cagcagaaat

20

&lt;210&gt; 16

&lt;211&gt; 20

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 16

ggtggctgtg gagacaaatc

20

35

&lt;210&gt; 17

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

<400> 17  
acttggacct ttccagcaac 20

<210> 18  
5<211> 19  
<212> DNA  
<213> Homo sapiens

<400> 18  
10ctttatccaa ccaggtgca 19

<210> 19  
<211> 23  
<212> DNA  
15<213> Homo sapiens

<400> 19  
tgggagaatt tagaaatgaa gga 23

20<210> 20  
<211> 21  
<212> DNA  
<213> Homo sapiens

25<400> 20  
tttcttcatt ttccctggtg a 21

<210> 21  
<211> 20  
30<212> DNA  
<213> Homo sapiens

<400> 21  
tggacagttt cccacattga 20  
35

<210> 22  
<211> 22  
<212> DNA  
<213> Homo sapiens  
40



&lt;400&gt; 22

ttcaaagggt gctgttctca aa

22

&lt;210&gt; 23

5&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 23

10tcaaacttct tgggcttaga aca

23

&lt;210&gt; 24

&lt;211&gt; 20

&lt;212&gt; DNA

15&lt;213&gt; Homo sapiens

&lt;400&gt; 24

cagagttgct ttcaatggca

20

20&lt;210&gt; 25

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25&lt;400&gt; 25

tccaggaaaa cttccttcca

20

&lt;210&gt; 26

&lt;211&gt; 18

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 26

ttcattggat acgtttcc

18

35

&lt;210&gt; 27

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

&lt;400&gt; 27

accagagttt cctgcaatgg

20

&lt;210&gt; 28

5&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 28

10tgccctgtgct gagtttgaat

20

&lt;210&gt; 29

&lt;211&gt; 20

&lt;212&gt; DNA

15&lt;213&gt; Homo sapiens

&lt;400&gt; 29

cggtcctcag tgtgcttgta

20

20&lt;210&gt; 30

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25&lt;400&gt; 30

ccaggatgag gactgggtaa

20

&lt;210&gt; 31

&lt;211&gt; 20

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 31

aagccgaaag gtgattgttg

20

35

&lt;210&gt; 32

&lt;211&gt; 23

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

<400> 32  
tatcatcttc attgtcctgc aga 23

<210> 33  
5<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 33  
10tcattgtcct gcagaagggtg 20

<210> 34  
<211> 20  
<212> DNA  
15<213> Homo sapiens

<400> 34  
gacgactcag aaaagccctg 20

20<210> 35  
<211> 20  
<212> DNA  
<213> Homo sapiens

25<400> 35  
aattggcagg aagcaacatc 20

<210> 36  
<211> 21  
30<212> DNA  
<213> Homo sapiens

<400> 36  
agtcagctcc tctgaacttt c 21  
35

<210> 37  
<211> 20  
<212> DNA  
<213> Homo sapiens  
40

<400> 37  
cgtgctcagc attctaacct 20

<210> 38  
5<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 38  
10gaacacctca ccttgtgcag 20

<210> 39  
<211> 21  
<212> DNA  
15<213> Homo sapiens

<400> 39  
cttgatagtc cagaaaaggc t 21

20<210> 40  
<211> 20  
<212> DNA  
<213> Homo sapiens

25<400> 40  
ccgcaagtct gtgcaataaa 20

<210> 41  
<211> 24  
30<212> DNA  
<213> Homo sapiens

<400> 41  
gtcagcttat gaagcctaatttct 24  
35

<210> 42  
<211> 22  
<212> DNA  
<213> Homo sapiens  
40

&lt;400&gt; 42

caaattgcac aggccctcta ga

22

&lt;210&gt; 43

5&lt;211&gt; 26

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 43

10aaagtctttt accctttcaa tagtca

26

&lt;210&gt; 44

&lt;211&gt; 23

&lt;212&gt; DNA

15&lt;213&gt; Homo sapiens

&lt;400&gt; 44

agagatttga gtttcaatgt ggg

23

20&lt;210&gt; 45

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25&lt;400&gt; 45

ctttaggctg gttgtcccaa

20

&lt;210&gt; 46

&lt;211&gt; 24

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 46

tgaaaactca ctcatttggt tcaa

24

35

&lt;210&gt; 47

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

&lt;400&gt; 47

cttcgagact ggacaagcca

20

&lt;210&gt; 48

5&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 48

10gagaggtcca ggaaggtaa

20

&lt;210&gt; 49

&lt;211&gt; 20

&lt;212&gt; DNA

15&lt;213&gt; Homo sapiens

&lt;400&gt; 49

acctggaggg agttcagaca

20

20&lt;210&gt; 50

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25&lt;400&gt; 50

aagagctgcc tctggtcctt

20

&lt;210&gt; 51

&lt;211&gt; 20

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 51

tacaagcaca ctgaggaccg

20

35

&lt;210&gt; 52

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

&lt;400&gt; 52

tttatgcagc cagcaagaag

20

&lt;210&gt; 53

5&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 53

10ggaggcaccc cttcttctaa

20

&lt;210&gt; 54

&lt;211&gt; 20

&lt;212&gt; DNA

15&lt;213&gt; Homo sapiens

&lt;400&gt; 54

gcggctctgg atgaagtgct

20

20&lt;210&gt; 55

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

25&lt;400&gt; 55

ctgagcaggg tcttctccac

20

&lt;210&gt; 56

&lt;211&gt; 20

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 56

agatgttgct tcctgccaat

20

35

&lt;210&gt; 57

&lt;211&gt; 20

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

<400> 57  
cagggccttt ctgagtcgtc 20

<210> 58  
5<211> 20  
<212> DNA  
<213> Homo sapiens

<400> 58  
10tgaacaagtg ttggacccag 20

<210> 59  
<211> 21  
<212> DNA  
15<213> Homo sapiens

<400> 59  
gattagcagc cctgcatatc t 21

20<210> 60  
<211> 20  
<212> DNA  
<213> Homo sapiens

25<400> 60  
gctcacagaa gcagtgagga 20

<210> 61  
<211> 20  
30<212> DNA  
<213> Homo sapiens

<400> 61  
taggctctga tatgccccat 20  
35

<210> 62  
<211> 10665  
<212> DNA  
<213> Homo sapiens  
40



&lt;220&gt;

&lt;221&gt; misc\_feature

&lt;222&gt; (1)...(10665)

&lt;223&gt; n = A,T,C or G

5

&lt;400&gt; 62

```

aaaatactcc cttgcctcaa aaactgctcg gtcaaacggt gatagcaaac cacgcattca      60
cagggccact gctgctcaca aaaccagtga ggatgatgcc aggatgatgt ctgcctcgcg      120
cctggctggg actctgatcc cagccatggc ctctctctcc tgcgtgagac cagaaagctg      180
10ggagccctgc gtggaggatg gtggctggag tcagctcttc tgaactttcc ctactttctg      240
cccagaactt ctactgtgtg gccctgggtt gtttattttt gcaaaaaaaaa aaagagttaa      300
attaccttaa agactcaaga agccacagag atcaaataat tcattgttac agggcactag      360
aggcagccat tgggggtttg ttccatttgg aaattttgag tgctaacagg ggcatgagat      420
aacatagatc tgcttaaggt cctgctctcg ctacctgtg gctctgtgaa gaaattatca      480
15aacctgtctg agactagttt tcgcatctgt aagagaatta taataccttc ttcactagag      540
agtaagcaga ctgcttcagt gtcatttctt cccactgggt gtctttacac tcagcttcaa      600
gcagtcaccc tgctcctttc aatctcagga aaaagatggc tttgtgtgtg tgtctctaga      660
gaaagaactt tctaagttgg tgcagacttc tgtatgcagt aatatagttt agtccagagg      720
atgaaaaaaaa taagagaatg aaaaaggaaa agagagagag agagaagaaa aaagcaagag      780
20ggaaatatgt ataatgtcag ctaatgcaac agtttctttc ttagtgaaat accaatcagc      840
tggttggtaa tcttattcat gatggatctc ttttgttttt cccctgcgca gacttcacag      900
ttgcttttaga aaccatagat agagccgaac agctaagaaa atgatttaca gtgaggcagg      960
gtcagaaaact caagagagaa aaagccagct gcagtcctga agttgaggat ataggagaaa     1020
atcaagtaat atttagcaaa gactaattca ttatcttgaa gccatccctt cctcaattc     1080
25cctgcccata gtctcctccc ttgtcctctt ctctgnatcc ctctgctgtt aggttaattg     1140
agatagattt tctaattang ctactgcca gataaaaccc agcccatgtt tctattagnc     1200
aatattgtct ttgaggctcc atggcttgca ncatttaagc agacatacga atgaagatct     1260
gcatgtttga actctgactt tgcgcattt acttcatttc tttgaatttc cattttcctc     1320
atctttaaat gcttatttga agattaagt aaagtatata acaaacaaga actatgcagg     1380
30cgtatggtaa gggattaatg atagatgata ataattaatg ttgacatcta ttgatcactt     1440
atactgtagc gggcttttaa ataaactctt taaacacctt atctcattta atccttcaaa     1500
cattctattg gtttcaaaca acagaaaact acaattagct ggcttctgca aggaattttg     1560
ttggaggaaa tgagagcatt cagaaattag atgggagcgt tagagaatta ggcttacaaa     1620
gaatgtggga aagtaggcta gaaagcagtg taaaaacaaa gacagcataa agcacttgac     1680
35cttatttact aggttcaccc atgggaatcc atgcactcta aagatttccc cctatttcta     1740
catcactttg ctcaagggtc aatgagccaa ggaaaagaat gcagttgtca aaatctgggc     1800
catgactaag gaaggtctgg acatcttgac tgccagacag tctcccaat gatatggagt     1860
atttgaatg atactggata ttttatttat tttttgtatt ttcaactttt aagttcagag     1920
gcacatgtgc agagcatgca ggtttattac ataagtaa atgtgtccatg gtgatttgct     1980
40gcatagatca tgaaaatatg gaacgcatca tggatttggt tgtcactcct gtgcaggggc     2040

```

catgctcacc	ttctctgtat	ccttccaatt	ttagtatatg	tgctactgca	gcaagcacga	2100
tattggatat	tttattacct	acattttaca	tatgataaaa	tgaggctcac	tgagggtttt	2160
cttttggtcg	ttttattttg	ttttgttttt	aaagacttgg	ccctaaacca	cacagaagag	2220
ctggcatgaa	acccagagct	ttcagactcc	ggagcctcag	cccttcaccc	cgattccatt	2280
5gcttcttget	aaatgctgcc	gttttatcnc	ggagggttaga	atgctgagca	cgtagtaggt	2340
gctctttact	ttctaactca	gagtaagaca	atttataagc	atgaattgag	tgaatggatg	2400
gatggatata	tgatggaag	gatggacaga	tgatgaaag	gttgactgaa	ttttgtgctt	2460
gcacaaaaag	aggccctct	ccaccatctc	tggtctagga	gaggggagtt	gggagaccat	2520
gcagtaaaag	tacttcatgt	catgtgtaat	cattgcaggt	ggttccta	attacttatt	2580
10aatgcatgga	gctgaatttc	tacaaaatcc	ccgacaacct	ccccttctca	accaagaacc	2640
tggaacctgag	ctttaatccc	ctgaggeatt	taggcagcta	tagcttcttc	agtttcccag	2700
aactgcaggt	gctggattta	tccaggtaat	gaatccactt	ttacatactg	cacaagggtga	2760
gggtgttcatt	gtctatcat	ttcattattg	gactggaaag	cttgggttgt	ggagtctcat	2820
cttcattcac	ttattcatte	atacaacaga	tgctttatta	actatataac	cttgagcaag	2880
15ctacctctat	tctccaggtc	tcagttttct	aatctgtgaa	gtaggcagtt	ggctgagaca	2940
gcttctaagg	gcaattctaa	ttttagggtt	tcttttaaga	caggagagaa	aattagctta	3000
aattctttca	taagcagcta	tttattgact	acttgcata	tggtgtacac	tctgcaagaa	3060
gacaggcata	tattgatata	taacacacag	cccctgttgt	taaggaggca	tatcttcttg	3120
aaagagttaa	taccttaaag	tcttgggtat	ggctctgggt	acatagtata	tagtcaacac	3180
20attttaatta	tgattttttg	gatctggaaa	ctgatataaa	gatagcgaca	tataacagta	3240
gggtgataaat	tatgttttaa	ctaaaggtaa	ctaattgtat	ttttcagaag	aggggccttc	3300
tctgtggtgg	gtagtcaaga	aagattcatg	aactgcataa	gattcaaaca	atgtctagaa	3360
tattaaaact	agtggtggca	ggtgaaatgt	catcttgata	ttttagggga	accaaattct	3420
aaaagggttt	tcatcatcgg	ggccttattt	gcaaatcgaa	ctagataatg	gatcatgttc	3480
25tctgcaatgg	tttgtaaaac	atttcaaaac	attttacata	ttttttatta	tagaaattat	3540
tgataaagac	taaggtcaca	gtataaaaat	ccttttttaga	gcagacattt	ctgtagaaga	3600
gtgaacatat	gacctattat	actctaattt	ggatatagat	aggatgtaac	aaaggagtaa	3660
tggaacaat	tcaaaggcag	tggtatagtg	catanagtcc	tggtggggtc	agaagacctg	3720
agcccaagtt	tacccccaac	atttataacc	catgtaacct	tagcatatta	cttcatctcc	3780
30cttaatcctt	agtttcatat	ctgatcaatg	gaaatgatga	aacttattct	gctggattaa	3840
atgtgataat	aaatattaat	atgctgtata	tatttaaatt	tttataaaat	atattttata	3900
agcataaagt	attcttacag	aatttcatta	ggtttttaaa	ataatttcaa	cttttatttt	3960
tgattcaggg	atttcatggt	ttatattgct	taatgctgag	gtgtagggta	caatcgatac	4020
catcactcag	gtagtgagca	tagtaccaa	tagttagttt	ttcaaccctt	gctgctttct	4080
35ctctatcccc	tctctagtaa	ccccagggt	ctatttttgt	catctttatg	tccatgtgta	4140
ctccatgttt	ggatcctact	tataaagtga	gaactcatgg	tatttggttt	tctgtncctt	4200
tggtngctaa	tttgcttagg	ataatggcta	ctagctgcat	ctatgccatt	atgttctaaa	4260
tttcanttnc	ctgcatgaaa	attttgtcaa	gtactctatt	aaggtagacc	acctctccct	4320
ttttttttca	aacaagaagt	agnttttccc	aaacaatgcc	cttatggaat	tnatcttcaa	4380
40tccnnggata	cccaataact	tgccccaaan	ccttaatctg	ncttacagag	aggccacctt	4440

ccttctgtaa cccataggag atttggattg gtaagaatgc tttgtgatag cccagcagcc 4500  
 ttctttcccc tatagaaata tatatatant ctttttatag gtgaggaact gaagcttgaa 4560  
 taatttaaata gacttatata catnattcatt gcttgttagc cacagaccag agatttaagt 4620  
 tcnatctccc agaatccaac ttaaattgtt tctttgtctt aatactctac ttctctaaag 4680  
 5tgattatcac caatgtaatg atatagagnc acagcaagac cctttccttc tcacctaattg 4740  
 tatagagcaa tgcagagata gaatgatggg ctataacaat catataattg aaagaaagaa 4800  
 cttcaaaaat aatcaagttc agctgtttga ttataaaatg tgataactaa aacctagaga 4860  
 ggaaaagagg tactcaagat cacacagtag gagaggactg cagaaacacc aaaccaagc 4920  
 tctttgtccc actcttccag cgttctttct actatactgc ctatccttta tctagtacc 4980  
 10aataaataac aaaagcttgg accacaatgc ttttattgtc taggaaactc ctgaagaagc 5040  
 taaataaaat ggggtgggaa tattgtaaat gtaattcagg ctggattaag aaagaactta 5100  
 tttgacattg taactgacaa gcacctgcaa tgctgaaagg aatttttcat tggcntgctg 5160  
 tttgtcgggc tgcataaag cctgtctctc aggacatgtc tctgaacatt gtgtgtagca 5220  
 tggccttcat ttcttttagg ataaaattca aaaccttta tctggttggg aaacctctgc 5280  
 15ctaattggga acctcttttc tccacaactc catattgtac actccaattt catctctgtt 5340  
 ctccaacct ggaagctatt tgcattgatt cctccttctg tcattttttt tctgtcaacc 5400  
 ttggggcttt tgtgtttgct gttcaactca cctcctttta ttgttaactt ctactcatct 5460  
 ttcaattttc aacttaagt ttctcagaga aacctacttt gattttcttg gtccanaacg 5520  
 gttctctgga tgtgaactct tatagcacat aattttcact tttttccaca aaactcgctc 5580  
 20ctatcacctg ttacaagcat ttacctctga taacaagaac tttcaaata ctagtgtca 5640  
 tgtaagcact ttccataaac attagagta tctgtgacac ttatgtgtaa tgtttcgtat 5700  
 ctctgaaatt gatatttacc agtcatttat ctggctacc aactaacaac tatccatatt 5760  
 atctgtacca atcagatgta taatcacaat ttgtgtgac agaaaatggc taaacttgat 5820  
 ccaaggctat tacatgcttt atcaactgca caatctttat atatgtcaat tattgatctt 5880  
 25taactgattt ccttcttatg gatcttctcc tctgcttate atgtatgcct aacatgacaa 5940  
 aaaagagcct atcattgcag ccagtatgat aatactcagt ctgtggggct tcttatttgc 6000  
 ttattccatc atcatctgtc ctgcttgatg tctttgcta tgcacaatca tatgacctat 6060  
 cacatctgta tgaagagctg gatgactagg attaatattc tatttttaggt tcttattcag 6120  
 cagaaatatt agataatcaa tgtcttttta ttctgtagg tgtgaaatcc agacaattga 6180  
 30agatggggca tatcagagcc taagccacct ctctacctta atattgacag gaaaccccat 6240  
 ccagagttaa gccctgggag ctttttctgg actatcaagt ttacagaagc tgggtggctgt 6300  
 ggagacaaat ctatcatctc tagagaactt cccattgga catctcaaaa ctttgaaaga 6360  
 acttaattgt gctcacaatc ttatccaatc ttcaaatta cctgagtatt tttctaactc 6420  
 gaccaatcta gagcacttgg acctttccag caacaagatt caaagtattt attgcacaga 6480  
 35cttgcggggt ctacatcaaa tgccctact caatctctct ttagacctgt ccctgaaccc 6540  
 tatgaacttt atccaaccag gtgcatttaa agaaattagg cttcataagc tgactttaag 6600  
 aaataatttt gatagtttaa atgtaatgaa aacttgtatt caaggctctg ctggtttaga 6660  
 agtccatcgt ttggttcttg gagaatttag aatgaagga aacttggaag agtttgacaa 6720  
 atctgctcta gagggcctgt gcaatttgac cattgaagaa tccccgatta gcatacttag 6780  
 40actactacct cgatgatatt attgacttat ttaattgggt gacaaatggg tcttcatttt 6840

ccctggtgag tgtgactatt gaaagggtaa aagacttttc ttataatttc ggatggcaac	6900
atttagaatt agttaactgt aaatttggac agtttccac attgaaactc aaatctctca	6960
aaaggcttac ttccacttcc aacaaagggtg ggaatgcttt ttcagaagtt gatctaccaa	7020
gccttgagtt tctagatctc agtagaaatg gcttgagttt caaagggtgc tgttctcaaa	7080
5gtgatttttg gacaaccagc ctaaagtatt tagatctgag cttcaatggg gttattacca	7140
tgagttcaaa cttcttgggc ttagaacaac tagaacatct ggatttccag cattccaatt	7200
tgaacaaat gagttagttt tcagtattcc tatcactcag aaacctcatt taccttgaca	7260
tttctcatac tcacaccaga gttgctttca atggcatctt caatggcttg tccagtctcg	7320
aagtcttgaa aatggctggc aattctttcc aggaaaactt ccttcagat atcttcacag	7380
10agctgagaaa cttgaccttc ctggacctct ctgagtgtca actggagcag ttgtctccaa	7440
cagcatttaa ctcactctcc agtcttcagg tactaaatat gagccacaac aacttctttt	7500
cattggatac gtttcttatt aagtgtctga actccctcca ggttcttgat tacagtctca	7560
atcacataat gacttccaaa aaacaggaac tacagcattt tccaagtagt ctgctttct	7620
taaactctac tcagaatgac ttgtcttgta cttgtgaaca ccagagtttc ctgcaatgga	7680
15tcaaggacca gaggcagctc ttgggtggaag ttgaacgaat ggaatgtgca acaccttcag	7740
ataagcaggg catgcctgtg ctgagtttga atatcacctg tcagatgaat aagaccatca	7800
ttgggtgtgc ggtccctcagt gtgctttag tatctgttgt agcagttctg gtctataagt	7860
tctattttca cctgatgctt cttgctggct gcataaagta tggtagaggt gaaaacatct	7920
atgatgcctt tgttatctac tcaagccagg atgaggactg ggtaaggaat gagctagtaa	7980
20agaatttaga agaaggggtg cctccatttc agctctgcct tcactacaga gactttattc	8040
ccggtgtggc cattgctgcc aacatcatcc atgaagggtt ccataaaagc cgaaagggtga	8100
ttgttgtggt gtcccagcac ttcattccaga gccgctgggt tatctttgaa tatgagattg	8160
ctcagacctg cgagtttctg agcagtcgtg ctggtatcat cttcattgtc ctgcagaagg	8220
tggagaagac cctgctcagg cagcaggtgg agctgtaccg ctttctcagc aggaacactt	8280
25acctggagtg ggaggacagt gtccctggggc ggcacatctt ctggagacga ctcagaaaag	8340
ccctgctgga tggtaaatca tggaaatccag aaggaacagt gggtagagga tgcaattggc	8400
aggaagcaac atctatctga agaggaaaa taaaaacctc ctgaggcatt tcttgcccag	8460
ctgggtccaa cacttgttca gttaataagt attaaatgct gccacatgtc aggccttatg	8520
ctaagggtga gtaattccat ggtgcactag atatgcaggg ctgctaattc caaggagctt	8580
30ccagtgcaga ggaataaat gctagactaa aatacagagt cttccagggtg ggcatttcaa	8640
ccaactcagt caaggaaacc atgacaaaaga aagtcatttc aactcttacc tcatcaagtt	8700
gaataaagac agagaaaaca gaaagagaca ttgttctttt cctgagtctt ttgaatggaa	8760
attgtattat gttatagcca tcataaaacc attttggtag ttttgactga actgggtgtt	8820
cactttttcc tttttgattg aatacaattt aaattctact tgatgactgc agtcgtcaag	8880
35gggctcctga tgcaagatgc ccttccatt ttaagtctgt ctccttacag aggttaaagt	8940
ctagtggcta attcctaagg aaacctgatt aacacatgct cacaaccatc ctggtcattc	9000
tcgagcatgt tctatttttt aactaatcac cctgatata tttttatttt tatatatcca	9060
gttttcattt ttttacgtct tgcctataag ctaatatcat aaataagggt gttaagacg	9120
tgttcaaat atccatatta accactatct ttcaaggaag tatggaaaag tacactctgt	9180
40cactttgtca ctcgatgtca ttccaaaggt attgcctact aagtaatgac tgtcatgaaa	9240

18

gcagcattga aataatttgt ttaaaggggg cactctttta aacgggaaga aaatttccgc 9300  
 ttcttgggtct tatcatggac aatttgggct ataggcatga aggaagtggg attacctcag 9360  
 gaagtcacct tttcttgatt ccagaaacat atgggctgat aaaccggggg tgacctcatg 9420  
 aaatgagttg cagcagatgt ttattttttt cagaacaagt gatgtttgat ggacctatga 9480  
 5atctatttag ggagacacag atggctggga tccctcccct gtacccttct cactgccagg 9540  
 agaactacgt gtgaagggtat tcaaggcagg gagtatacat tgctgtttcc tgttgggcaa 9600  
 tgctccttga ccacattttg ggaagagtgg atgttatcat tgagaaaaca atgtgtctgg 9660  
 aattaatggg gttcttataa agaagggtcc cagaaaagaa tgttcattcc agcttcttca 9720  
 ggaaacagga acattcaagg aaaaggacaa tcaggatgtc atcagggaaa tgaaaataaa 9780  
 10aaccacaatg agatatcacc ttataccagg tagatggcta ctataaaaa atgaagtgtc 9840  
 atcaaggata tagagaaatt ggaacccttc ttcactgctg gagggaatgg aaaatggtgt 9900  
 agccgttatg aaaaacagta cggagggttc tcaaaaatta aaaatagaac tgctatatga 9960  
 tccagcaatc tcacttctgt atatatacc aaaataattg aaatcagaat ttcaagaaaa 10020  
 tatttacact cccatgttca ttgtggcact cttcacaatc actgtttcca aagttatgga 10080  
 15aacaacccaa atttccattg gaaaataaat ggacaaagga aatgtgcata taacgtacaa 10140  
 tggggatatt attcagccta aaaaaagggg ggatcctgtt atttatgaca acatgaataa 10200  
 acccgaggc cattatgcta tgtaaaatga gcaagtaaca gaaagacaaa tactgcctga 10260  
 tttcatttat atgaggttct aaaatagtca aactcataga agcagagaat agaacagtgg 10320  
 ttctagggga aaaggaggaa gggagaaatg aggaaatagg gagttgtcta attggtataa 10380  
 20aattatagta tgcaagatga attagctcta aagatcagct gtatagcaga gttcgtataa 10440  
 tgaacaatac tgtattatgc acttaacatt ttgttaagag ggtacctctc atgttaagtg 10500  
 ttcttaccat atacatatac acaagggaagc ttttggaggt gatggatata tttattacct 10560  
 tgattgtggg gatggtttga caggatgtg actatgtcta aactcatcaa attgtataca 10620  
 ttaaatatat gcagttttat aatatcaaaa aaaaaaaaaa aaaaa 10665

25

&lt;210&gt; 63

&lt;211&gt; 28

30&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 63

attttgtttt gtttttaaag acttggcc

28

35

&lt;210&gt; 64

&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

40

&lt;400&gt; 64

gtcatgtgta atcattgcag gtggttcc

28

&lt;210&gt; 65

5&lt;211&gt; 28

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 65

10atgtcttttt attcctgtag gtgtgaaa

28

&lt;210&gt; 66

&lt;211&gt; 30

&lt;212&gt; DNA

15&lt;213&gt; Artificial Sequence

&lt;220&gt;

&lt;223&gt; A primer

20&lt;400&gt; 66

gattagcata cttagactac tacctccatg

30

&lt;210&gt; 67

&lt;211&gt; 27

25&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 67

gatcaacttc tgaaaaagca ttccac

27

30

&lt;210&gt; 68

&lt;211&gt; 31

&lt;212&gt; DNA

&lt;213&gt; Artificial Sequence

35

&lt;220&gt;

&lt;223&gt; A primer

&lt;400&gt; 68

40ggttgctgtt ctcaaagtga ttttgggaga a

31

<210> 69  
 <211> 30  
 <212> DNA  
 <213> Homo sapiens

5

<400> 69  
 acctgaagac tggagagtga gttaaatgct

30

<210> 70  
 10<211> 1360  
 <212> DNA  
 <213> Homo sapiens

<220>  
 15<221> misc\_feature  
 <222> (1)...(1360)  
 <223> n = A,T,C or G

<400> 70  
 20ttccacttct aagagctgcc tagagtagtc aagattatag agacaaaagt agtgcataga 60  
 ttcaggggcc tagggaaaagg ggaaatgggg agttatttat taatgaatag tggatgatgat 120  
 tgtacaaaaa tatgaacata attaatgcc aataattgtn cacatacaaa tggatcaagat 180  
 aataaatttt atgttatgtc atgttatgtt atgtgatattt accataatac agaaaatgaa 240  
 aaaagaaaag aaagaaagta aagcttagcg gtttncatga cttgnccaat gcctcaaagc 300  
 25catgagtcga cccagctgag atctganctt cagtatatcc cattctgaaa tcccagactt 360  
 tcccgaatct tcttgacttt ttcaaactgt gtttcagttg aggtttattt tcagttttgt 420  
 atgtgagttt cttcgcaaga agggcgggcc aaattgtgtc ctgcaaaaac ctacatatcg 480  
 aagtcctaac ccctctacct cagactatga ctgtatatgg agagagagcc ttgaaagagg 540  
 tatgtaaggt agaattgagg cattatgggt ggcctaatc caacataact ggtgtcctta 600  
 30taagaagggg agattagaat tcagacacac ttgtgacac cttgagttca gactggaagc 660  
 ctctagaatt gtgagaaaat gaatgtctgt tgtttaagcc acccagtcctg tggatatttc 720  
 ttatggcagc cccagcaaac taatacaaat agtgtttcca cagctgaaac aaaattggaa 780  
 aatcacctgc atcctagaga gttacaaggg ctattttaat agaacctgat tgttttccta 840  
 aattcaccaa gccagggcag aggtcagatg actaatggg ataaaagcca actagcttcc 900  
 35tcttgctgtt tcttttagcca ctggctctgca ggcgttttct tcttctaact tcctctcctg 960  
 tgacaaaaga gataactatt agagaaacaa aagtcagaa tgctaagggt gccgctttca 1020  
 ctctctctca ccctttagcc cagaactgct ttgaatacac caattgctgt ggggcggctc 1080  
 gaggaagaga agacaccagt gcctcagaaa ctgctcgggt agacgggtgat agcgagccac 1140  
 gcattcacag ggccactgct gctcacagaa gcagtggaga tgatgccagg atgatgtctg 1200  
 40cctcgcgcct ggctgggact ctgatccag ccatggcctt cctctcctgc gtgagaccag 1260

21

aaagctggga gccctgcgtg gaggtatgtg gctggagtca gctcctctga actttccctc 1320  
acttctgccc agaacttctc actgtgtgcc ctggtttgtt 1360

&lt;210&gt; 71

5&lt;211&gt; 1333

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;400&gt; 71

10cgcatcatgg atttgtgtgt catccttgtg caggggcat gctcatcttc tctgtatcct 60  
tccaatttta gtatatgtgc tactgcagca agcacgatat tggatatattt attacctaca 120  
ttttacatat gataaaatga ggctcactga ggtttttctt ttgttcgttt tattttgttt 180  
tgtttttaaa gacttggccc taaaccacac agaagagctg gcatgaaacc cagagctttc 240  
agactccgga gcctcagccc ttcaccccga ttcatttgct tcttgctaaa tgctgccgtt 300  
15ttatcacgga ggtagaatg ctgagcacgt agtaggtgct ctttactttc taatctagag 360  
taagacaatt tataagcatg aattgagtga atggatggat ggatatatgg atggaaggat 420  
ggacagatgg atgaaagggt gactgaattt tgtgcttgca caaaaagagg cccctctcca 480  
ccatctctgg tctaggagag gggagttggg agaccatgca gtaaagatac ttcattgcat 540  
gtgtaatcat tgcaggtggg tcctaataat acttatcaat gcatggagct gaatttctac 600  
20aaaatccccg acaacctccc cttctcaacc aagaacctgg acctgagctt taatccccctg 660  
aggcatttag gcagctatag cttcttcagt tccccagaac tgcaggtgct ggatttatcc 720  
aggtaatgaa tccactttta catactgcac aaggtagggt gttcattgtc ctatcatttc 780  
attattggac tggaaagctt ggtttgtgga gtctcatctt cattcactta ttcattcata 840  
caacagatgt cttattaact atataacctt gagcaagcta cctctattct ccaggtctca 900  
25gttttctaat ctgtgaagta ggcagttggc tgagacagct tctaagggca attctaattt 960  
taggttttct ttttaagacag gagagaaaat tagcttaaatt tctttcataa gcagctattt 1020  
attgactact tgctatatgt tgtacactct gcaagaagac aggcataat tgatatataa 1080  
cacacagccc ctgttggtta ggaggcatac cttcttgaaa gagttaatac cttaaagtcc 1140  
tggttatggc cctgggtaca tagtatatag tcaacacatt ttaattatga ttttttgat 1200  
30ctggaaactg atataaagat agcgacatat aacagtaggt gataaattat gtttaacta 1260  
aaggtaacta attgtatttt tcagaagagg ggccttctct gtggtgggta gtcaagaaag 1320  
attcatgaac tgc 1333

&lt;210&gt; 72

35&lt;211&gt; 6786

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

40&lt;221&gt; misc\_feature



&lt;222&gt; (1) ... (6786)

&lt;223&gt; n = A,T,C or G

&lt;400&gt; 72

```

5ggtaagaatg ctttgtgata gccacgcagc cttctttccc ctatagaaat atatatan 60
  tctttttata ggtgaggaac tgaagcttga ataatttaaa tgacttatat acatnatcat 120
  tgcttgtagg ccacagacca gagatttaag ttcncatctc cagaatccaa cttaaatggt 180
  ttctttgtct taatactcta cttctctaaa gtgattatca ccaatgtaat gatatagagn 240
  cacagcaaga ccctttccct ctcacctaat gtatagagca atgcagagat agaatgatgg 300
10gctataacaa tcatataatt gaaagaaaga acttcaaaaa taatcaagtt cagctgtttg 360
  atttataaat gtgataacta aaacctagag aggaaaagag gtactcaaga tcacacagta 420
  ggagaggact gcagaaacac caaacccaag ctcttttgtc cactcttcca gcgttctttc 480
  tactatactg cctatccctt atctagttac caataataaa caaaagcttg gaccacaatg 540
  cttttattgt ctaggaaact cctgaagaag ctaaataaaa tgggtgggga atattgtaaa 600
15tgtaattcag gctggattaa gaaagaactt atttgacatt gtaactgaca agcacctgca 660
  atgctgaaag gaatttttca ttggcntgct gtttgctggg ctgcatcaaa gccctgtctc 720
  taggacatgt ctctgaacat tgtgtgtagc atggctttca tttcttttag gataaaatc 780
  aaaacccttt atctggttgg taaacctctg cctaattggg aaccttcttt ctccacaact 840
  ccatattgta cactccaatt tcatctctgt tctccaacca tggaagctat ttgtcatgat 900
20tctctcttgt gtcatttttt ttctgtcaac cttggggctt ttgtgtttgc tgttcaactc 960
  acctcttttt attgttaact tctactcatc tttcaatttt caacttaagt gttctcagag 1020
  aaacctactt tgattttctt ggtccanaac gggtctctgg atgtgaactc ttatagcaca 1080
  taattttcac ttttttccac aaaactcgct cctatcacct gttacaagca tttacctctg 1140
  ataacaagaa ctttcaataa tctagctgtc atgtaagcac ttttcataaa cattaagagt 1200
25atctgtgaca cttatgtgta atgtttcgta tctctgaaat tgatatttac cagtcattta 1260
  tcttggttac caactaaca ctatccatat tatctgtacc aatcagatgt ataatacaca 1320
  ttttgtgtga cagaaaatgg ctaaacttga tccaaggcta ctacatgctt tatcaactgc 1380
  acaatcttta tatatgtcaa ttattgatct ttanctgatt tcttctttat ggattttctc 1440
  ctctgcttat catgtatgcc taacatgaca aaaaagagcc tatcattgca gccagtatga 1500
30taataactcag tctgtggggc ttcttatttg cttattccat catcatctgt cctgcttgat 1560
  gtctttgcct atgcacaatc atatgacca tcacatctgt atgaagagct ggatgactag 1620
  gattaatatt ctatttttagg ttcttattca gcagaaatat tagataatca atgtcttttt 1680
  attcctgtag gtgtgaaatc cagacaattg aagatggggc atatcagagc ctaagccacc 1740
  tctctacctt aatattgaca ggaaaccca tccagagttt agccctggga gccttttctg 1800
35gactatcaag ttacagaag ctggtggctg tggagacaaa tctagcatct ctagagaact 1860
  tccccattgg acatctcaaa actttgaaag aacttaatgt ggctcacaat cttatccaat 1920
  ctttcaaatt acctgagtat ttttctaata tgaccaatct agagcacttg gacctttcca 1980
  gcaacaagat tcaaagtatt tattgcacag acttgcgggg tctacatcaa atgcccctac 2040
  tcaatctctc tttagacctg tccctgaacc ctatgaactt tatccaacca ggtgcattta 2100
40aagaaattag gcttcataag ctgactttaa gaaataattt tgatagttaa aatgtaatga 2160

```

aaacttgat	tcaaggtctg	gctggtttag	aagtccatcg	tttggttctg	ggagaattta	2220
gaaatgaagg	aaacttgga	aagtttgaca	aatctgctct	agagggcctg	tgcaatttga	2280
ccattgaaga	attccgatta	gcatacttag	actactacct	cgatgatatt	attgacttat	2340
ttaattgttt	gacaaatgtt	tcttcatttt	ccctgggtgag	tgtgactatt	gaaagggtaa	2400
5aagacttttc	ttataatttc	ggatggcaac	atrtagaatt	agttaactgt	aaatttggac	2460
agtttccac	attgaaactc	aaatctctca	aaaggcttac	tttacttcc	aacaaagggtg	2520
ggaatgcttt	ttcagaagtt	gatctacca	gccttgagtt	tctagatctc	agtagaaatg	2580
gcttgagttt	caaagggtgc	tgttctcaaa	gtgattttgg	gacaaccagc	ctaaagtatt	2640
tagatctgag	cttcaatggt	gttattacca	tgagttcaaa	cttcttgggc	ttagaacaac	2700
10tagaacatct	ggatttccag	cattccaatt	tgaaacaaat	gagtgaagttt	tcagtattcc	2760
tactactcag	aaacctcatt	taccttgaca	tttctcatat	tcacaccaga	gttgctttca	2820
atggcatctt	caatggcttg	tccagtctcg	aagtcttgaa	aatggctggc	aattctttcc	2880
aggaaaactt	ccttccagat	atcttccag	agctgagaaa	cttgaccttc	ctggacctct	2940
ctcagtgtca	actggagcag	ttgtctccaa	cagcatttaa	ctcactctcc	agtcttcagg	3000
15tactaaatat	gagccacaac	aacttctttt	cattggatac	gtttcttat	aagtgtctga	3060
actcctcca	ggttcttgat	tacagtctca	atcacataat	gacttccaaa	aaacaggaac	3120
tacagcattt	tccaagtagt	ctagctttct	taaatcttac	tcagaatgac	tttgcttgta	3180
cttgatgaaca	ccagagtttc	ctgcaatgga	tcaaggacca	gaggcagctc	ttggtggaag	3240
ttgaacgaat	ggaatgtgca	acaccttcag	ataagcaggg	catgctgtg	ctgagtttga	3300
20atatacctg	tcagatgaat	aagaccatca	ttggtgtgtc	ggctctcagt	gtgcttgtag	3360
tatctgttgt	agcagttctg	gtctataagt	tctattttca	cctgatgctt	cttgctggct	3420
gcataaagta	tggtagaggt	gaaaacatct	atgatgcctt	tggtatctac	tcaagccagg	3480
atgaggactg	ggtaagggaat	gagctagtaa	agaatttaga	agaaggggtg	cctccatttc	3540
agctctgcct	tactacaga	gactttattc	ccgggtgtggc	cattgctgcc	aacatcatcc	3600
25atgaagggtt	ccataaaagc	cgaaagggtga	ttgttgtggt	gtcccagcac	ttcatccaga	3660
gccgtgggtg	tatctttgaa	tatgagattg	ctcagacctg	gcagtttctg	agcagtcgtg	3720
ctggtatcat	cttcattgtc	ctgcagaagg	tggagaagac	cctgctcagg	cagcaggtgg	3780
agctgtaccg	ccttctcagc	aggaacactt	acctggagtg	ggaggacagt	gtcctggggc	3840
ggcacatctt	ctggagacga	ctcagaaaag	ccctgctgga	tggtaaatca	tggaaatccag	3900
30aaggaaacagt	gggtacagga	tgcaattggc	aggaagcaac	atctatctga	agaggaaaaa	3960
taaaaacctc	ctgaggcatt	tcttgcccag	ctgggtccaa	cacttgttca	gttaataagt	4020
attaatatgct	gccacatgtc	aggccttatg	ctaagggtga	gtaattccat	ggtgcactag	4080
atatgcaggg	ctgctaattct	caaggagctt	ccagtgcaga	gggaataaat	gctagactaa	4140
aatacagagt	cttccaggtg	ggcattttcaa	ccaactcagt	caaggaaccc	atgacaaaga	4200
35aagtcatttc	aactcttacc	tcatcaagtt	gaataaagac	agagaaaaca	gaaagagaca	4260
ttgttctttt	cctgagtcct	ttgaatggaa	attgtattat	gttatagcca	tcataaaacc	4320
attttggtag	ttttgactga	actgggtgtt	cactttttcc	tttttgattg	aatacaattt	4380
aaattctact	tgatgactgc	agtcgtcaag	gggctcctga	tgcaagatgc	cccttccatt	4440
ttaagtctgt	ctccttacag	aggttaaagt	ctagtggcta	attcctaagg	aaacctgatt	4500
40aacacatgct	cacaaccatc	ctggctcattc	tcgagcatgt	tctatttttt	aactaatcac	4560

ccctgatata	tttttatttt	tatatateca	gttttcattt	ttttacgtct	tgcctataag	4620
ctaataatcat	aaataagggt	gtttaagacg	tgcttcaaatt	atccatatta	accactattt	4680
ttcaaggaag	tatggaaaag	tacactctgt	cactttgtca	ctcgatgtca	ttccaaaggt	4740
attgcctact	aagtaatgac	tgatcatgaaa	gcagcattga	aataatttgt	ttaaagggg	4800
5cactctttta	aacgggaaga	aaatttccgc	ttcttgggt	tatcatggac	aatttgggct	4860
agaggcagga	aggaagtggg	atgacctcag	gaggtcacct	tttcttgatt	ccagaaacac	4920
atgggctgat	aaacccgggg	tgacctcatg	aaatgagttg	cagcagaagt	ttattttttt	4980
cagaacaagt	gatgtttgat	ggacctctga	atctcttttag	ggagacacag	atggctggga	5040
tcctctccct	gtaccttctt	cactgccagg	agaactacgt	gtgaagggtat	tcaaggcagg	5100
10gagtatacat	tgtctgtttcc	tggtgggcaa	tgctccttga	ccacattttg	ggaagagtgg	5160
atgttatcat	tgagaaaaca	atgtgtctgg	aattaatggg	gttcttataa	agaagggttcc	5220
cagaaaagaa	tgttcatcca	gcctcctcag	aaacagaaca	ttcaagaaaa	ggacaatcag	5280
gatgtcatca	gggaaatgaa	aataaaaacc	acaatgagat	atcaccttat	accaggtaga	5340
atggctacta	taaaaaaatg	aagtgtcatc	aaggatatag	agaaattgga	acccttcttc	5400
15actgctggag	ggaatggaaa	atgggtgtagc	cgttatgaaa	aacagtacgg	agggtttctca	5460
aaaattaaaa	atagaactgc	tatatgatcc	agcaatctca	cttctgtata	tatacccaaa	5520
ataattgaaa	tcagaatttc	aagaaaatat	ttacactccc	atgttcattg	tggcactctt	5580
cacaatcact	gtttccaaag	ttatggaaac	aacccaaatt	tccattgaaa	aataaatgga	5640
caaagaaaat	gtgcatatac	gtacaatggg	atattattca	gcctaaaaaa	agggggnatc	5700
20ctgttattta	tgacaacatg	aataaaccgc	gagccattat	gctatgtaaa	atgagcaagt	5760
aacagaaaga	caaatactgc	ctgatttcat	ttatatgagg	ttctaaaata	gtcaaaactca	5820
tagaagcaga	gaatagaaca	gtggttccta	gggaaaagga	ggaagggaga	aatgaggaaa	5880
tagggagttg	tctaattggg	ataaaattat	agtatgcaag	atgaattagc	tctaaagatc	5940
agctgtatag	cagagtctgt	ataatgaaca	atactgtatt	atgcacttaa	cattttgtta	6000
25agaggggtacc	tctcatgtta	agtgttctta	ccatatacat	atacacaagg	aagcttttgg	6060
agggtgatga	tatatattat	accttgattg	tggtgatggg	ttgacaggta	tgtgactatg	6120
tctaaactca	tcaaatgtga	tacattaaat	atatgcagtt	ttataatatc	aattatgtct	6180
gaatgaagct	ataaaaaaga	aaagacaaca	aaattcagtt	gtcaaaaactg	gaaatatgac	6240
cacagtcaga	agtgtttgtt	actgagtggt	tcagagtggt	tttgggttga	gcagggtctag	6300
30ggtgattgaa	catccctggg	tgtgtttcca	tgtctcatgt	actagtgaag	gtagatgtgt	6360
gcatttgtgc	acatatccct	atgtatccct	atcagggtcg	tggttatattg	aaagtgtgtg	6420
tgtccgcag	atcatatctg	tatagaagag	agtgtgatta	tatttcttga	agaatacatc	6480
catttgaaat	ggatgtctat	ggctgtttga	gatgagttct	ctactcttgt	gcttgtacag	6540
tagtctcccc	ttatccctta	tgcttgggtg	atacgttctt	agaccccaag	tggatctctg	6600
35agaccgcaga	tggtaccaaa	cctcatatat	gcaatatttt	ttctatatac	taaataccta	6660
agataaagtt	catcttctga	attaggcaca	gtaagagatt	aacaataact	aacaataaaa	6720
ttgaatagtt	ataataatat	attgtaataa	aagttatgtg	aatgtgatct	ctttcttttc	6780
tctctc						6786







FIG. 1

2/18



3/18





FIG. 3A



SUBSTITUTE SHEET (RULE 26)

4/18



5/18

↓

Human (aa 290)	.	.	.	L	A	Y	L	D	Y	Y	L	D	D	I	I	D	L	F	N	C	L	T	N	V	.	.	.
Mouse (aa 289)	.	.	.	L	T	Y	T	N	D	F	S	D	D	I	V	K	-	F	H	C	L	A	N	V	.	.	.
Rat (aa 289)	.	.	.	L	T	Y	I	N	H	F	S	D	D	I	Y	N	-	L	N	C	L	A	N	I	.	.	.
Hamster (aa 289)	.	.	.	F	T	Y	A	N	E	F	S	E	D	I	T	D	-	F	D	C	L	A	N	V	.	.	.

FIG. 4



6/18




FIG. 5

7/18




FIG. 6A

8/18




FIG. 6B

9/18




FIG. 6C

10/18




FIG. 6D

11/18

<u>Amino Acid</u>	<u>Codon</u>
Phe	UUU, UUC
Ser	UCU, UCC, UCA, UCG, AGU, AGC
Tyr	UAU, UAC
Cys	UGU, UGC
Leu	UUA, UUG, CUU, CUC, CUA, CUG
Trp	UGG
Pro	CCU, CCC, CCA, CCG
His	CAU, CAC
Arg	CGU, CGC, CGA, CGG, AGA, AGG
Gln	CAA, CAG
Ile	AUU, AUC, AUA
Thr	ACU, ACC, ACA, ACG
Asn	AAU, AAC
Lys	AAA, AAG
Met	AUG
Val	GUU, GUC, GUA, GUG
Ala	GCU, GCC, GCA, GCG
Asp	GAU, GAC
Gly	GGU, GCC, GGA, GGG
Glu	GAA, GAG

**FIG. 7**

12/18

Original Residue	Exemplary Substitutions	Preferred Substitutions
Ala (A)	val; leu; ile	val
Arg (R)	lys; gln; asn	lys
Asn (N)	gln; his; lys; arg	gln
Asp (D)	glu	glu
Cys (C)	ser	ser
Gln (Q)	asn	asn
Glu (E)	asp	asp
Gly (G)	pro	pro
His (H)	asn; gln; lys; arg	arg
Ile (I)	leu; val; met; ala; phe norleucine	leu
Leu (L)	norleucine; ile; val; met; ala; phe	ile
Lys (K)	arg; gln; asn	arg
Met (M)	leu; phe; ile	leu
Phe (F)	leu; val; ile; ala	leu
Pro (P)	gly	gly
Ser (S)	thr	thr
Thr (T)	ser	ser
Trp (W)	tyr	tyr
Tyr (Y)	trp; phe; thr; ser	phe
Val (V)	ile; leu; met; phe; ala; norleucine	leu

FIG. 8

SUBSTITUTE SHEET (RULE 26)

13/18

## HUMAN TLR4 GENOMIC SEQUENCE

```

AAAATACTCC CTTGCCTCAA AAAGTGTCTG GTCAAACGGT
GATAGCAAAC CACGCATTCA CAGGGCCACT GCTGCTCACA
AAACCAGTGA GGATGATGCC AGGATGATGT CTGCCTCGCG
CCTGGCTGGG ACTCTGATCC CAGCCATGGC CTTCTCTCTC
TGCGTGAGAC CAGAAAGCTG GGAGCCCTGC GTGGAGGTAT
GTGGCTGGAG TCAGCTCCTC TGAACCTTCC CTCACCTCTG
CCCAGAAGTT CTCACCTGTG GCCCTGGTTT GTTTATTTT
GCAAAAAAAA AAAGAGTTAA ATTACCTTAA AGACTCAAGA
AGCCACAGAG ATCAAATAAT TCATTGTTAC AGGGCACTAG
AGGCAGCCAT TGGGGGTTTG TTCCATTTGG AAATTTTGAG
TGCTAACAGG GGCATGAGAT AACATAGATC TGCTTAAGGT
CCCTGCTCTG CTACCTTGTG GCTCTGTGAA GAAATTATCA
AACCTGTCTG AGACTAGTTT TCGCATCTGT AAGAGAATTA
TAATACCTTC TTCACTAGAG AGTAAGCAGA CTGCTTCAGT
GTCATTTCTT CCCACTGGTG GTCTTTACAC TCAGCTTCAA
GCAGTCACCC TGCTCCTTTC AATCTCAGGA AAAAGATGGC
TTTGTGTGTG TGTCTCT:A: G:AGAAAGAA CTTTCTAAGT
TGGTGCAGA CTTCTGTATG CAGTAATATA GTTTAGTCCA
GAGGATGAAA AAAATAAGAG A:ATGAAAAA GGAAAAGAGA
GAGAGAGA:G AAGAAAAAAG CAAGAGGGAA AT:ATGTATA
ATGTCAGCTA ATGCAAC:AG TTTCTTTCTT AGTGAAATAC
CAATCAGCTG :GTTG:GTAA TCTT:ATTCA TGATGGATCT
CTTTTGT TTTT TCCCCTGCGC AGACTTC:AC AGTTGCTTTA
GAAACCCATA GTAGAGCCGA A:CAGCTAAG AAAATGATTT
ACAGTGAGGC AGGGTCAGAA ACTCAAGAGA GAAAAAGCCA
GCTGCAGTC: CTGAAGT:TG AGGATATAGG :AGAAAAATCA
AGTAATATTT AGCAAAGACT AATTCATTAT CTTGAAGCCA
TCCCTTCCCT CAATTCCCTG CCCATAGTCC TCCTCCTTGT
CCTCTTCTCT GNA:TCCCTC TGCTGTTAGG TTA:ATGG:A
GATAGATTTT CTAATTANGC TCACTGCGAG ATAAAACCCA
GCCCATGTTT CTATTAGNCA ATATTGTCTT TGAGGCTCCA
TGGCTTGCAN CATTTAAGCA GACATACGAA TGAAGATCTG
CATGTTTGAA CTCTGACTTT GCGCATATTA CTTCAATTCT
TTGAATTTCC ATTTTCCTCA TCTTTAAATG CTTATTTGAA
GATTAAGTGA AAGTATATAA CAAACAAGAA CTATGCAGGC
GTATGGTAAG GGATTAATGA TAGATGATAA TAATTAATGT
TGACATCTAT TGATCACTTA TACTGTAGCG GGCTTTTAAA
TAAACTCTTT AAACACCTTA TCTCATTTAA TCCTTCAAAC
ATTCTATTGG TTTCAAACAA CAGAAAACTA CAATTAGCTG
GCTTCTGCAA GGAATTTTGT TGGAGGAAAT GAGAGCATTC
AGAAATTAGA TGGGAGCGTT AGAGAATTAG GCTTACAAAG
AATGTGGGAA AGTAGGCTAG AAAGCAGTGT AAAAACAAAG
ACAGCATAAA GCACTTGACC TTATTTACTA GGTTCACCA
TGGGAATCCA TGCACCTTAA AGATTTCCTT CTATTTCTAC
ATCACTTTGC TCAAGGGTCA ATGAGCCAAG GAAAAGAATG

```

FIG. 9

SUBSTITUTE SHEET (RULE 26)



14/18

CAGTTGTCAA AATCTGGGCC ATGACTAAGG AAGGTCTGGA  
CATCTTGACT GCCAGACAGT CTCCCCAATG ATATGGAGTA  
TTTAGAATGA TACTGGATAT TTTATTTATT TTTTGTATTT  
TCAACTTTTA AGTTCAGAGG CACATGTGCA GAGCATGCAG  
GTTTATTACA TAAGTAAATG TGTGCCATGG TGATTTGCTG  
CATAGATCAT GAAAATATSG AACGCATCAT GGATTTGTGT  
GTCATCCTTG TGCAGGGGCC ATGCTCATCT TCTCTGTATC  
CTTCCAATTT TAGTATATST GCTACTGCAG CAAGCACGAT  
ATTGGATATT TTATTACCTA CATTTTACAT ATGATAAAAT  
GAGGCTCACT GAGGTTTTTC TTTTGTTCTG TTTATTTTGT  
TTTGTTTTTTA AAGACTTGGC CCTAAACCAC ACAGAAGAGC  
TGGCATGAAA CCCAGAGCTT TCAGACTCCG GAGCCTCAGC  
CCTTCACCCC GATTCCATTG CTTCTTGCTA AATGCTGCCG  
TTTTATCNCG GAGGTTAGAA TGCTGAGCAC GTAGTAGGTG  
CTCTTTACTT TCTAATCTAG AGTAAGACAA TTTATAAGCA  
TGAATTGAGT GAATGGATGG ATGGATATAT GGATGGAAGG  
ATGGACAGAT GGATGAAAGG TTGACTGAAT TTTGTGCTTG  
CACAAAAGA GGCCCTCTC CACCATCTCT GGTCTAGGAG  
AGGGGAGTTG GGAGACCATG CAGTAAAGAT ACTTCATGTC  
ATGTGTAATC ATTGCAGGTG GTTCCTAATA TTACTTATCA  
ATGCATGGAG CTGAATTTCT ACAAATCCC CGACAACCTC  
CCCTTCTCAA CCAAGAACCT GGACCTGAGC TTTAATCCCC  
TGAGGCATTT AGGCAGCTAT AGCTTCTTCA GTTTCCCAGA  
ACTGCAGGTG CTGGATTTAT CCAGGTAATG AATCCACTTT  
TACATACTGC ACAAGGTGAG GTGTTCAATG TCCTATCATT  
TCATTATTGG ACTGGAAAGC TTGGTTTGTG GAGTCTCATC  
TTCATTCACT TATTCATTCA TACAACAGAT GTCTTATTAA  
CTATATAACC TTGAGCAAGC TACCTCTATT CTCCAGGTCT  
CAGTTTTCTA ATCTGTGAAG TAGGCAGTTG GCTGAGACAG  
CTTCTAAGGG CAATTCTAAT TTTAGGTTTT CTTTTAAGAC  
AGGAGAGAAA ATTAGCTTAA ATTCTTTCAT AAGCAGCTAT  
TTATTGACTA CTTGCTATAT GTTGTAACAT CTGCAAGAAG  
ACAGGCATAT ATTGATATAT AACACACAGC CCCTGTTGTT  
AAGGAGGCAT ATCTTCTTGA AAGAGTTAAT ACCTTAAAGT  
CCTGGGTATG GTCCTGGGTA CATAGTATAT AGTCAACACA  
TTTTAATTAĀ GATTTTTTGG ATCTGGAAAC TGATATAAAG  
ATAGCGACAT ATAACAGTAG GTGATAAATT ATGTTTAAAC  
TAAAGGTAAC TAATTGTATT TTTCAGAAGA GGGGCCTTCT  
CTGTGGTGGG TAGTCAAGAA AGATTCATGA ACTGCATAAG  
ATTCAAACAA TGTCTAGAAT ATTAAAACTA GTGGTGGCAG  
GTGAAATGTC ATCTTGATAT TTTAGGGGAA CCAAATTCTA  
AAAGGGTTTT CATCATCGGG GCCTTATTTG CAAATCGAAC  
TAGATAATGG ATCATGTTCT CTGCAATGGT TTGTAAAACA  
TTTCAAAACA TTTTACATAT TTTTATTAT AGAAATTATT  
GATAAAGACT AAGGTCACAG TATAAAAATC CTTTTTAGAG  
CAGACATTTT TGTAGAAGAG TGAACATATG ACCTATTATA  
CTCTAATTTG GATATAGATA GGATGTAACA AAGGAGTAAT

FIG. 9 (Continued)

SUBSTITUTE SHEET (RULE 26)

15/18

GGGAACAAT<sup>1</sup> CAAAGGCAGT GGTATAGTGC ATANAGTCCT  
GTTGGGGTCA GAAGACCTGA GCCCAAGTTT ACCCCCAACA  
TTTATAACCC ATGTAACCTT AGCATATTAC TTCATCTCCC  
TTAATCCTTA GTTTCATATC TGATCAATGG AAATGATGAA  
ACTTATTCTG CTGGATTAAA TGTGATAATA AATATTAATA  
TGCTGTATAT ATTTAAATTT TTATAAAATA TATTTTATAA  
GCATAAAGTA TTCTTACAGA ATTTCATTAG GTTTTTAAAA  
TAATTTCAAC TTTTATTTTT GATTACAGGA TTTACATGGT  
TATATTGCGT AATGCTGAGG TGTAGGGTAC AATCGATACC  
ATCACTCAGG TAGTGAGCAT AGTACCCAAT AGTTAGTTTT  
TCAACCCCTG CTGCTTTCTC TCTATCCCCT CTCTAGTAAT  
CCCCAGGGTC TATTTTTGTC ATCTTTATGT CCATGTGTAC  
TCCATGTTTG GATCCTACTT ATAAAGTGAG AACTCATGGT  
ATTTGGCTTT CTGTNCCTTT GTTNGCTAAT TTGCTTAGGA  
TAATGGCTAC TAGCTGCATC TATGCCATTA TGTCTAAAT  
TTCANTNCC TGCATGAAAA TTTTGTCAAG TACTCTATTA  
AGGTAGACCA CCTCTCCCTT TTTTTTCAA ACAAGAAGTA  
GNTTTTCCCA AACAATGCCC TTATGGAATT NATCTTCAAT  
CCNNGGATAC CCAATAACTT GCCCAAANC CTTAATCTGN  
CTTACAGAGA GGCCACCTTC CTTCTGTAAC CCATAGGAGA  
TTTGGATTGG TAAGAATGCT TTGTGATAGC CCAGCAGCCT  
TCTTTCCCCT ATAGAAATAT ATATATANTC TTTTATAGG  
TGAGGAAC<sup>2</sup> AAGCTTGAAT AATTTAAATG ACTTATATAC  
ATNATCAT<sup>3</sup> CTGTGTTAGC ACAGACCAGA GATTTAAGTT  
CNCATCTCCA GAATCCA<sup>4</sup>ACT TAAATGTTTT CTTTGTCTTA  
ATACTCTACT TCTCTAAAGT GATTATCACC AATGTAATGA  
TATAGAGNCA CAGCAAGACC CTTTCCTTCT CACCTAATGT  
ATAGAGCAAT GCAGAGATAG AATGATGGGC TATAACAATC  
ATATAATTGA AAGAAAGAAC TTCAAAAATA ATCAAGTTCA  
GCTGTTTGAT TTATAAATGT GATAACTAAA ACCTAGAGAG  
GAAAAGAGGT ACTCAAGATC ACACAGTAGG AGAGGACTGC  
AGAAACACCA AACCCAAGCT CTTTGTGCCA CTCTCCAGC  
GTTCTTTCTA CTATACTGCC TATCCTTTAT CTAGTTACCA  
ATAAATAACA AAAGCTTGGA CCACAATGCT TTTATTGTCT  
AGGAAACTCC TGAAGAAGCT AAATAAAATG GGTGGGGAAT  
ATTGTAAATG TAATTCAGGC TGGATTAAGA AAGAACTTAT  
TTGACATTGT AACTGACAAG CACCTGCAAT GCTGAAAGGA  
ATTTTTCATT GGCNTGCTGT TTGCTGGGCT GCATCAAAGC  
CCTGTCTCTA GGACATGTCT CTGAACATTG TGTGTAGCAT  
GGCTTTCATT TCTTTTAGGA TAAAATTCAA AACCCTTTAT  
CTGGTTGGTA AACCTCTGCC TAATTGGGAA CCTTCTTTCT  
CCACAAC<sup>5</sup>TCC ATATTGTACA CTCCAATTC ATCTCTGTTC  
TCCAACCATG GAAGCTATTT GTCATGATTC CTCCTTGTGT  
CATTTTTTTT CTGTCAACCT TGGGGCTTTT GTGTTTGCTG  
TTCACTTCAC CTCCTTTTAT TGTAACTTC TACTCATCTT  
TCAATTTTCA ACTTAAGTGT TCTCAGAGAA ACCTACTTTG  
ATTTTCTTGG TCCANAACGG TTCTCTGGAT GTGAACTCTT

FIG. 9 (Continued)

SUBSTITUTE SHEET (RULE 26)

16/18

ATAGCACATA ATTTTCACTT TTTTCCACAA AACTCGCTCC  
 TATCACCTGT TACAAGCATT TACCTCTGAT AACAAGAAGT  
 TTCAAATATC TAGCTGTCAT GTAAGCACTT TTCATAAACA  
 TTAAGAGTAT CTGTGACACT TATGTGTAAT GTTTCGTATC  
 TCTGAAATTG ATATTTACCA GTCATTTATC TTGGCTACCA  
 ACTAACAAGT ATCCATATTA TCTGTACCAA TCAGATGTAT  
 AATCACAATT TTGTGTGACA GAAAATGGCT AAACCTTGATC  
 CAAGGCTATT ACATGCTTT: ATCAACTGCA CAATCTTTAT  
 ATATGTCAAT TATTGATCTT TAACTGATTT CCTTCTTATG  
 :GATTTTCTC CTCTGCTTAT CATGTATGCC TAACAT:GAC  
 AAAAAAG:AG CCTA:TCATT GCAGCCAGTA TGATAATACT  
 CA:GTCTGTG GGGCTTCTTA TTTGCTTAT: TCCATCATCA  
 TCTGTCTGCTG TTGATGTCTT TGCCTATGCA CAATCATATG  
 :ACCCATCAC ATCTGTATGA AGAGC:TGGA TGAAGTAGGAT  
 TAATATTCT: AT::TTTAG GTTCTTATT: CAGCAGAAAT  
 ATTAGATAA: TCAATGTCTT TTTATTCCTG TAGGTGTGAA  
 ATCCAGACAA TTGAAGATGG GGCATATCAG AGCCT:AAGC  
 CACCTCTCTA CCTTAATATT GACAGGAAAC CCCATCCAGA  
 GTTTAGCCCT GGGAGCCCTT TCTGGACTAT CAAGTTTACA  
 GAAGCTGGTG GCTGTGGAGA CAAATCTAGC ATCTCTAGAG  
 AACTTCCCCA TTGGACATCT CAAAACCTTG AAAGAACTTA  
 ATGTGGCTCA CAATCTTATC CAATCTTTCA AATTACCTGA  
 GTATTTTCT AATCTGACCA ATCTAGAGCA CTGGGACCTT  
 TCCAGCAACA AGATTCAAAG TATTTATTGC ACAGACTTGC  
 GGGTTCTACA TCAAATGCC CTACTCAATC TCTCTTTAGA  
 CCTGTCCCTG AACCCATATGA ACTTTATCCA ACCAGGTGCA  
 TTTAAAGAAA TTAGGCTTCA TAAGCTGACT TTAAGAAATA  
 ATTTTGATAG TTTAAATGTA ATGAAAACCT GTATTCAAGG  
 TCTGGCTGGT TTAGAAGTCC ATCGTTTGGT TCTGGGAGAA  
 TTTAGAAATG AAGGAACTT GGAAAAGTTT GACAAATCTG  
 CTCTAGAGGG CCTGTGCAAT TTGACCATTG AAGAATTCCC  
 GATTAGCATA CTTAGACTAC TACCTCGATG ATATTATTGA  
 CTTATTTAAT TGTTTGACAA ATGGTTCTTC ATTTTCCCTG  
 GTGAGTGTTGA CTATTGAAAG GGTAAGGAC TTTTCTTATA  
 ATTTCCGGATG GCAACATTTA GAATTAGTTA ACTGTAAATT  
 TGGACAGTTT CCCACATTGA AACTCAAATC TCTCAAAGG  
 CTTACTTTCA CTTCCAACAA AGGTGGGAAT GCTTTTTCAG  
 AAGTTGATCT ACCAAGCCTT GAGTTTCTAG ATCTCAGTAG  
 AAATGGCTTG AGTTTCAAAG GTTGCTGTTT TCAAAGTGAT  
 TTTGGGACAA CCA:GCCT:A AAGTATTTAG ATCTGAGCTT  
 CAATGGTGTT A:TTACCATG AGTTCAAAC TCTTGGGCTT  
 AGAACA:ACT AGAACATCTG GATTTCAGC ATTCCAATTT  
 GAAACA:AAT GAGTGAGTTT TCAGTATTC TA:TCACCA  
 GAAA:CCT:C ATTTACCTTG ACATTTCTCA TACTCACACC  
 AGAGTTGCTT TCAATGGCAT CTTCAATGGC TTGTCCAGTC  
 TCGAAGTCTT GAAAATGGCT GGCAATTCTT TCCAGGAAAA  
 CTTCTTCCA GATATCTTCA CAGAGCTGAG AAACCTTGACC

FIG. 9 (Continued)

SUBSTITUTE SHEET (RULE 26)

17/18

TTCCTGGACC TCTCTCAGTG TCAACTGGAG CAGTTGTCTC  
CAACAGCATT TAACCTCACTC TCCAGTCTTC AGGTACTAAA  
TATGAGCCAC AACCAACTTCT TTTCATTGGA TACGTTTCCT  
TATAAGTGTC TGAACCTCCCT CCAGGTTCTT GATTACAGTC  
TCAATCACAT AATGACTTCC AAAAAACAGG AACTACAGCA  
TTTTCCAAGT AGTCTAGCTT TCTTAAATCT TACTCAGAAT  
GACTTTGCTT GTACTTGTGA ACACCAGAGT TTCCTGCAAT  
GGATCAAGGA CCAGAGGCAG CTCTTGGTGG AAGTTGAACG  
AATGGAATGT GCAACACCTT CAGATAAGCA GGGCATGCCT  
GTGCTGAGTT TGAATATCAC CTGTCAGATG AATAAGACCA  
TCATTGGTGT GTCGGTCCTC AGTGTGCTTG TAGTATCTGT  
TGTAGCAGTT CTGGTCTATA AGTTCTATTT TCACCTGATG  
CTTCTTGCTG GCTGCATAAA GTATGGTAGA GGTGAAAAACA  
TCTATGATGC CTTTGTTATC TACTCAAGCC AGGATGAGGA  
CTGGGTAAGG AATGAGCTAG TAAAGAATTT AGAAGAAGGG  
GTGCCTCCAT TTCAGCTCTG CCTTCACTAC AGAGACTTTA  
TTCCCGGTGT GGCCATTGCT GCCAACATCA TCCATGAAGG  
TTTCCATAAA AGCCGAAAGG TGATTGTTGT GGTGTCCCAG  
CACTTCATCC AGAGCCGCTG GTGTATCTTT GAATATGAGA  
TTGCTCAGAC CTGGCAGTTT CTGAGCAGTC GTGCTGGTAT  
CATCTTCATT GTCCTGCAGA AGGTGGAGAA GACCCTGCTC  
AGGCAGCAGG TGGAGCTGTA CCGCCTTCTC AGCAGGAACA  
CTTACCTGGA GTGGGAGGAC AGTGTCTTGG GGCGGCACAT  
CTTCTGGAGA CGACTCAGAA AAGCCCTGCT GGATGGTAAA  
TCATGGAATC CAGAAGGAAC AGTGGGTACA GGATGCAATT  
GGCAGGAAGC AACATCTATC TGAAGAGGAA AAATAAAAAAC  
CTCCTGAGGC ATTTCTTGCC CAGCTGGGTC CAACACTTGT  
TCAGTTAATA AGTATTAAAT GCTGCCACAT GTCAGGCCTT  
ATGCTAAGGG TGAGTAATTC CATGGTGCAC TAGATATGCA  
GGGCTGCTAA TCTCAAGGAG CTTCCAGTGC AGAGGGAATA  
AATGCTAGAC TAAAATACAG AGTCTTCCAG GTGGGCATTT  
CAACCAACTC AGTCAAGGAA CCCATGACAA AGAAAGTCAT  
TTCAACTCTT ACCTCATCAA GTTGAATAAA GACAGAGAAA  
ACAGAAAGAG ACATTGTTCT TTTCCTGAGT CTTTTGAATG  
GAAATTGTAT TATGTTATAG CCATCATAAA ACCATTTTGG  
TAGTTTTGAC TGAACGGGT GTTCACTTTT TCCTTTTGA  
TTGAATACAA TTTAAATTCT ACTTGATGAC TGCAGTCGTC  
AAGGGGCTCC TGATGCAAGA TGCCCCCTCC ATTTTAAGTC  
TGTCTCCTTA CAGAGGTTAA AGTCTAGTGG CTAATTCCTA  
AGGAAACCTG ATTAACACAT GCTCACAACC ATCCTGGTCA  
TTCTCGAGCA TGTTCTATTT TTTAACTAAT CACCCCTGAT  
ATATTTTAT TTTATATAT CCAGTTTTCA TTTTTTTACG  
TCTTGCTTAT AAGCTAATAT CATAAATAAG GTTGTTTAAG  
ACGTGCTTCA AATATCCATA TTAACCACTA TTTTCAAGG  
AAGTATGGAA AAGTACACTC TGTCACCTTG TCACTCGATG  
TCATTCCAAA GTTATTGCCT ACTAAGTAAT GACTGTCATG  
AAAGCAGCAT TGAAATAATT TGTTTAAAGG GGGCACTCTT

FIG. 9 (Continued)

SUBSTITUTE SHEET (RULE 26)

18/18

```

TTAAACGGGA AGAAAATTTT CGCTTCCTGG TCTTATCATG
GACAAATTTGG GCTATAGGCA TGAAGGAAGT GGGATTACCT
CAGGAAGTCA CCTTTTCITG ATTCCAGAAA CATATGGGCT
GATAAACCCG GGGTGACCTC ATGAAATGAG TTGCAGCAGA
TGTTTATTTT TTTCAGAACA AGTGATGTTT GATGGACCTA
TGAATCTATT TAGGGAGACA CAGATGGCTG GGATCCCTCC
CCTGTACCCT TCTCACTGCC AGGAGAACTA CGTGTGAAGG
TATTCAAGGC AGGGAGTATA CATTGCTGTT TCCTGTTGGG
CAATGCTCCT TGACCACATT TTGGGAAGAG TGGATGTTAT
CATTGAGAAA ACAATGTGTC TGAATTAAT GGGGTTCTTA
TAAAGAAGGT TCCCAGAAAA GAATGTTTCT TCCAGCTTCT
TCAGGAAACA GGAACATTCA AGGAAAAGGA CAATCAGGAT
GTCATCAGGG AAATGAAAAT AAAAACCACA ATGAGATATC
ACCTTATACC AGGTAGATGG CTAATAATAA AAAATGAAGT
GTCATCAAGG ATATAGAGAA ATTGGAACCC TTCTTCACTG
CTGGAGGGAA TGGAAAATGG TGTAGCCGTT ATGAAAAACA
GTACGGAGGT TTCTCAAAAA TTAAAAATAG AACTGCTATA
TGATCCAGCA ATCTCACTTC TGTATATATA CCAAAAATAA
TTGAAATCAG AATTTCAAGA AAATATTTAC ACTCCCATGT
TCATTGTGGC ACTCTTCACA ATCACTGTTT CCAAAGTTAT
GGAAACAACC CAAATTTCCA TTGGAAAAATA AATGGACAAA
GGAAATGTGC ATATAACGTA CAATGGGGAT ATTATTCAGC
CTAAAAAAG GGGGATCCT GTTATTTATG ACAACATGAA
TAAACCCGGA GGCCATTATG CTATGTAAAA TGAGCAAGTA
ACAGAAAGAC AAATACTGCC TGATTTTATT TATATGAGGT
TCTAAAATAG TCAAACTCAT AGAAGCAGAG AATAGAACAG
TGGTTCCTAG GGAAAAGGAG GAAGGGAGAA ATGAGGAAAT
AGGGAGTTGT CTAATTGGTA TAAAATTATA GTATGCAAGA
TGAATTAGCT CTAAAGATCA GCTGTATAGC AGAGTTCGTA
TAATGAACAA TACTGTATTA TGCCTTAAC ATTTTGTAA
GAGGGTACCT CTCATGTTAA GTGTTCTTAC CATATACATA
TACACAAGGA AGCTTTTGGA GGTGATGGAT ATATTTATTA
CCTTGATTGT GGTGATGGTT TGACAGGTAT GTGACTATGT
CTAACTCAT CAAATTGTAT ACATTAAATA TATGCAGTTT
TATAATATCA AAAAAAAAAA AAAAAAAAAA

```

FIG. 9 (Continued)

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/15723

A. CLASSIFICATION OF SUBJECT MATTER  
 IPC 7 C12N15/12 C12N15/63 C12N5/10 C12Q1/68 C07K14/705  
 G01N33/68 A01K67/027

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
 IPC 7 C12N C12Q C07K G01N A01K

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, BIOSIS, CHEM ABS Data, MEDLINE

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	<p>ALEXANDER POLTORAK ET AL.: "Defective LPS signaling in C3H/HeJ and C57BL/10ScCr mice: Mutations in Tlr4 gene" SCIENCE, vol. 282, 11 December 1998 (1998-12-11), pages 2085-2088, XP002145980 LANCASTER, PA US cited in the application abstract page 2086, right-hand column, last paragraph -page 2087, left-hand column, paragraph 2; figure 2</p> <p style="text-align: center;">--- -/--</p>	<p>1-3,5,6, 8,9, 11-21, 23,24, 28-30</p>

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

## \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*B\* document member of the same patent family

Date of the actual completion of the international search

19 September 2000

Date of mailing of the international search report

04/10/2000

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentaan 2  
 NL - 2280 HV Rijswijk  
 Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
 Fax: (+31-70) 340-3016

Authorized officer

Montero Lopez, B

## INTERNATIONAL SEARCH REPORT

International Application No

PCT/US 00/15723

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SALMAN T. QURESHI ET AL.: "Endotoxin-tolerant mice have mutations in Toll-like receptor (Tlr4)" THE JOURNAL OF EXPERIMENTAL MEDICINE, vol. 189, no. 4, 15 February 1999 (1999-02-15), pages 615-625, XP000900609 cited in the application page 616, left-hand column, paragraph 3 page 618, right-hand column, paragraph 2 -page 623, right-hand column, last paragraph -----	1-3,5,6, 8,9, 11-21, 23,24, 28-30
X	WO 98 50547 A (SCHERING CORPORATION) 12 November 1998 (1998-11-12)  page 7, line 5 - line 35 page 8, line 26 -page 10, line 12 page 12, line 21 -page 13, line 11 page 14, line 32 -page 18, line 9 page 24, line 19 - line 30 page 29, line 23 -page 30, line 23 page 37, line 20 -page 43, line 11 page 52, line 33 -page 53, line 4 -----	1,2,5,8, 11-20, 23,28-30
P,X	N.C. ARBOUR ET AL.: "Missense mutations of the TLR4 gene are associated with hyporesponsiveness to lipopolysaccharides (LPS) in humans" AMERICAN JOURNAL OF HUMAN GENETICS, vol. 65, no. 4, October 1999 (1999-10), page A97 XP000900485 abstract no. 506 -----	1-24, 28-32
P,X	NANCY C. ARBOUR ET AL.: "TLR4 mutations are associated with endotoxin hyporesponsiveness in humans" NATURE GENETICS, vol. 25, no. 2, June 2000 (2000-06), pages 187-191, XP000900719 the whole document -----	1-24, 28-35

## FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

Continuation of Box I.2

Claims Nos.: 25-27

Present claims 25-27 relate to an agent defined by reference to a desirable characteristic or property, namely altering TLR4 activity. The claims cover all agents having this characteristic or property, whereas the application does not provide support within the meaning of Article 6 PCT and/or disclosure within the meaning of Article 5 PCT for any specific example of such agent. In the present case, the claims so lack support, and the application so lacks disclosure, that a meaningful search over the whole of the claimed scope is impossible. Independent of the above reasoning, the claims also lack clarity (Article 6 PCT). An attempt is made to define the agent by reference to a result to be achieved. Again, this lack of clarity in the present case is such as to render a meaningful search over the whole of the claimed scope impossible. Consequently, no search has been carried out for claims 25-27.

The applicant's attention is drawn to the fact that claims, or parts of claims, relating to inventions in respect of which no international search report has been established need not be the subject of an international preliminary examination (Rule 66.1(e) PCT). The applicant is advised that the EPO policy when acting as an International Preliminary Examining Authority is normally not to carry out a preliminary examination on matter which has not been searched. This is the case irrespective of whether or not the claims are amended following receipt of the search report or during any Chapter II procedure.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

national Application No

PCT/US 00/15723

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO 9850547 A	12-11-1998	AU 7175498 A	27-11-1998
		EP 0980429 A	23-02-2000
		NO 995458 A	08-11-1999
<hr/>			